

**STUDIES ON THE EFFECT OF SOME
PESTICIDES ON BLOOD PARAMETERS IN FRESH
WATER FISHES OF BUNDELKHAND REGION**



Thesis submitted to
Bundelkhand University, Jhansi
for the degree of
Doctor of philosophy
in
Zoology

(November 2009)



Under guidance of
Dr. Kaneez Zahra
Sr. Lecturer

Submitted by
Saurabh Shreshth
M.Sc. (Zoology)

**DEPARTMENT OF ZOOLOGY
BIPIN BEHARI (P.G.) COLLEGE,
BUNDELKHAND UNIVERSITY, JHANSI, U.P.**

Time swallows up

Everything that is visible,

Sparing nothing,

It does not spare even

Outstanding personalities.



*This Thesis is Dedicated in words and
spirits to my revered Grand Father
Late Shri Maiyadeen*

Dr. Kaneez Zahra

(Senior lecturer)

Department of Zoology

Bipin Bihari P.G. College

Jhansi 284001

Ref.

Date4/11/09

Certificate

This is to certify that the thesis entitled "**Studies on the effect of some pesticides on blood parameters in fresh water fishes of Bundelkhand region**" embodies the original research work of **Saurabh Shreshth** who worked under my supervision and guidance for more than five years in the department of Zoology, Bipin Bihari (P.G.) College, Jhansi.

This thesis has not been submitted for any degree to any other university.



Dr. Kaneez Zahra

(Senior lecturer)

Department of Zoology

Bipin Bihari P.G. College

Jhansi 284001

CONTENTS

Acknowledgement	I-III
Chapter 01	
Introduction	1-13
Chapter 02	
Review of literature	14-25
Chapter 03	
Experimental design and Methodology	26-39
01. Haemoglobin percentage	
02. TEC	
03. TLC	
04. ESR	
05. MCH	
06. MCHC	
07. MCV	
09. Blood glucose	
Chapter 04	
(a.) Chemistry of toxicant	40-49
01. Glyphosate	
02. Phosphamidon	
03. Metasystox	
04. Imidacloprid	
(b.) Physiochemical properties of water	49
Chapter 05	
Effect of Glyphosate on blood parameters of <i>Channa punctatus</i>	50-72
01. Result	

- (a.) Acute toxicity bioassay and behavior
- (b.) Haematological & Biochemical study (Acute & Chronic)
- (c.) Study of seasonal variation

02. Discussion

- (a.) Acute toxicity bioassay and behavior
- (b.) Haematological & Biochemical study
- (c.) Study of seasonal variation

Chapter 06

Effect of Phosphamidon on blood parameters of *Channa punctatus* 73-96

01. Result

- (a.) Acute toxicity bioassay and behavior
- (b.) Haematological & Biochemical study (Acute & Chronic)
- (c.) Study of seasonal variation

02. Discussion

- (a.) Acute toxicity bioassay and behavior
- (b.) Haematological & Biochemical study
- (c.) Study of seasonal variation

Chapter 07

Effect of Metasystox on blood parameters of *Channa punctatus* 97-121

01. Result

- (a.) Acute toxicity bioassay and behavior
- (b.) Haematological & Biochemical study (Acute & Chronic)
- (c.) Study of seasonal variation

02. Discussion

- (a.) Acute toxicity bioassay and behavior
- (b.) Haematological & Biochemical study
- (c.) Study of seasonal variation

Chapter 08

Effect of Imidacloprid on blood parameters of *Channa punctatus* 122-144

01. Result

- (a.) Acute toxicity bioassay and behavior
- (b.) Haematological & Biochemical study (Acute & Chronic)
- (c.) Study of seasonal variation

02. Discussion

- (a.) Acute toxicity bioassay and behavior
- (b.) Haematological & Biochemical study
- (c.) Study of seasonal variation

Summery 145-151

Bibliography 152-189

Aberrations and symbols

ACKNOWLEDGEMENT

It gives me mammoth fulfillment in expressing and indebtedness towards my supervisor **Dr. (Mrs.) Kaneez Zahra**, Senior lecturer, Department of Zoology, Bipin Bihari (P.G.) College, Jhansi for her precious direction, steady encouragement and kind help during the entire tenure of my research work. She is a supervisor not only formal sense but intrinsically a devoted personality in full capacity. I have no expression words to illustrate her innumerable qualities of mind and heart.

My obligation and honest thanks to the respectable principal **Dr. S.N. Shrivastava**, Bipin Bihari College Jhansi, I am extremely obliged to him.

I express my heartiest thanks to **Dr. A. B. Gupta** head of the department in Zoology, Bipin Bihari (P.G.) College, Jhansi for providing essential laboratory facilities for this research work. Emphatically owe my sincere thanks to **Dr. V. I. Sharma** for their constructive interference.

No expression of thanks will be sufficient with recognition of intelligence and professional dexterity of **Dr. A. K. Shrivastav**, **Dr. O.P. Yadav**, **Dr. S. K Dube**, **Dr. Fatima Siddique** and **Dr. Hemant kumar B.B.C.** Jhansi for having meticulously support my work with unflinching help and suggestions at every stage of its preparation.

Financial assistance was one of the main factors and I owe profound gratitude to **University Grants Commission, New Delhi** for their support to me in the form of research fellowship.

Credit goes also to **Dr. Anil Kumar** department in Zoology, Bipin Bihari (P.G.) College, Jhansi for providing informations in chemistry.

I extend my sincere credit to all the other research scholars friends **Mr. Jag Mohan Sen, Mr. Ashish Kumar Vishwkarma, Mr. Raj Kumar Rai, Smt Sahin Naz** etc.

I gratefully acknowledge all the departmental non teaching staff of Zoology (**Shri Mukesh kumar, Shri Mahendra, Shri Bhawani Prasad, Shri Mushtaq Khan, Shri Ram dayal, Shri Rakesh Tripathy and Shri Mahendra Kumar**) for inspirations and providing help in care and managing of experiments.

Mr. Sharad Kumar contributes to the completion of practical work such as use, handling, care of fishes and proper maintenance of experiments.

I emphatically owe my sincere thanks to **Director ITRC, Lucknow** for their constant help of library from time to time.

It is my pleasure to convey my thanks to **Shri Ramlal Kushwaha, Shri Ram Gopal Verma (L.T.) and Shri Ram Swaroop Kushwaha.**

I wish to offer my sincere gratitude and regards by heart to my respected **Father Shri Ramkishun, Mother Smt Kalawati** who gave me best possible education, honesty, support, affection, blessing and inspiration to make me utmost from next to nothing. I have no expressions to convey my gratefulness to my elder brother **Er. Harikesh Kumar Shreshth, Bhahi Smt. Santoshi Shreshth, Sister Km. Saumya Shreshth** for helping me directly and indirectly during whole work.



(Saurabh Shreshth)

Chapter - 1

INTRODUCTION

INTRODUCTION

The study of Toxicology has generated greater excitement and become a subject of general interest in the present century. Hippocrates was the first to introduce primitive principals of toxicology around 400 BC. More significant contributions of Toxicology were made in 16th century and Paracelsus (the father of toxicology) was the first to introduce the wide theory of Toxicology. Toxicology is derived from the word toxicum means poison, and logus means knowledge i.e. “the science of poison”. It can be defined as, “A discipline of science which deals with the various aspects of poison and poisoning is known as Toxicology”. Poison may be defined as a substance which even in small dose produces adverse effects in metabolism of an organism and consequently may cause death.

Toxicology is the study of how specific chemicals cause injury to living cells and organism. It is a study to determine how easily the chemical enters the organism, what cells are affected by the chemicals and what cell functions are impaired. Toxicological study is becoming an increasingly important subject of modern society because of the adverse effects of the growing use of chemicals and high-tech radioactive appliances. With increasing population, the modern society demands improvement of the health and living conditions such as nutrition, clothing, dwelling and transportation. In order to fulfill these growing needs a large variety of chemicals must be manufactured and used. These chemical compounds come in contact with various segments of the population (animal and human beings) directly or indirectly causing adverse effect on them.

Toxicology is multi disciplinary science. It includes Environmental, Economic, Clinical and Forensic toxicology. The environmental degradation due to the presence of various pollutants is known as environmental toxicology. My research topic is related to the environmental toxicology. Environmental toxicology is normally based on the pollution studies. In general pollution can be defined as a matter in wrong place or any substance released into the environment which degrades it. Our environment becomes continuously polluted due to continued economic growth, mismanagement of resources, population growth and more use of toxicants (Stout B.A., 1976; Kutty et al., 1977; Suege J. and Bluzat R., 1983; Kulshrestha S.K. and Arora A., 1984; PAN Pesticide Database 1996; Kumar Suresh et al., 1999). A toxicant may be defined as an agent that causes adverse effect or response in biological system, seriously damaging its structure or function or producing death. The adverse effect or response may be defined in the term of a management that is out side the normal range for healthy organism. The toxicants are released from various sources into air, water and soil. They get into human food chain from the environment. Once the toxicants enter our biological system, they interfere with the biochemical process as may lead to fatal results.

Aquatic Toxicology: -

A large number of toxicants reach in the aquatic environment. The sources of toxicants in aquatic environment are :-

- Sewage and other wastes of domestic nature
- Agricultural discharges

- Industrial effluents
- Wastes from thermal and nuclear power plants

Aquatic toxicants may be grouped into two categories

1. Toxic trace elements and heavy metals found in natural and wastes water,
2. Pesticides

Classification of pesticides:-

A pest is described as the trouble some or destructive organism and pesticides may be defined as substances intended for killing pest. Pesticides may be classified in various ways

1. Classification of pesticides by target organism:-

- Acaricides : Used against mites and ticks
- Algaecide : Used against algae
- Fungicides : Used against fungi
- Molluscicide : Used to kill molluscs pest and other invertebrates
- Nematicide : Used against Nematodes
- Herbicides : Weed kills
- Insecticides : Used against insects
- Rodenticides : Used against mice, rats and other rodents

2. Classification on the basis of chemical classes:-

On the basis of chemical nature, the pesticides may be broadly divided into two groups

a. Organic pesticides

b. Inorganic pesticides

At present, the uses of inorganic pesticides are less than organic pesticides. Mostly the synthetic organic pesticides are being used in developed and developing countries. These are organochlorine, organophosphorus, natural and synthetic pyrethroide and dinitrophenols etc.

Contamination by pesticides

There are various ways through which the pesticides add into the environment either by direct or indirect applications (Heath AG., 1995; Heath AG., 1996; Braunbeck, T., 1994; Mason C.F. ,1996; Hrudey, S.E., W. Chen and C.G. Roussex, 1996).

(A) Direct application of pesticides .

The pesticides are directly applied to water bodies for any of the following purposes;

- (i) for the control of unwanted weeds.
- (ii) for the control of insect pests infecting water plants.
- (iii) for the control of insects and their life-stages of public-health importance, for instance mosquito larvae.

- (iv) for the control of undesired fish in fish culture ponds to restock with more desirable fish.

(B) Indirect application of pesticides

The pesticides indirectly reach the aquatic reservoirs by any of the following ways:

- (i) A heavy rainfall soon after the pesticide application in agricultural fields washes away fairly large share of the pesticides with the run off water, otherwise only a small share of pesticides reach into water bodies through agricultural run off. The water soluble pesticides are transported in dissolved state while insoluble pesticides get bound to particulate matter, which is carried by water.
- (ii) Pesticides manufacturing factories may release large amounts of pesticides, even in the effluents, which are treated for the removal of pesticides.
- (iii) Pesticides may also reach water- bodies on account of fall out from accidental spray from large scale aerial spraying of forest or agricultural field's. Usually major fall out occurs close to the site of their application polluting nearby water- bodies, but some times wind may carry the drift to considerable distances.


Production and Usage of pesticide in India

The word pesticide covers a broad range of compounds including insecticides, herbicides, pesticides, plant growth regulators and others. The production of pesticides started in India in 1952 with

the establishment of a plant for the production of organophosphorus pesticides near Calcutta, and India is now the second largest manufacturer of pesticides in Asia after China and ranks twelfth globally. There has been a steady growth in the production of pesticides in India, from 80,000 metric tones; in 1988 to 102,240 metric tones in 1998. In 2008 it became to 134,004 metric tones. The use of herbicides is correspondingly less. The most important crops with regards to insecticidal uses are cotton, fruits, vegetables, cereal and maize.

General information of toxicants:-

Several organophosphorus insecticides have been reviewed by WHO for consideration as agents for the control of disease vectors. A large number has been reviewed by the FAO/WHO Joint Meetings on Pesticide Residues. These compounds are designed to be toxic for certain pests and are added deliberately to the environment. Organophosphorus insecticides are normally esters, amides, or thiol derivatives of phosphoric, phosphonic, phosphorothioic, or phosphonothioic acids. The organophosphorus pesticides are derivatives of phosphoric acid having one or more, rarely two atoms of phosphorus whose one or more hydrogen atoms are placed by alkyl groups. General formula of organophosphorus compound is



where R¹ and R² are usually simple alkyl or aryl groups, both of which may be bonded directly to phosphorus (in phosphonates), or

linked via -O-, or -S- (in phosphates), or R1 may be bonded directly and R2, and X bonded via one of the above groups (phosphonates).

The primary biochemical effect associated with toxicity caused by organophosphorus pesticides is inhibition of AChE. The normal function of AChE is to terminate neurotransmission due to ACh that has been liberated at cholinergic nerve endings in response to nervous stimuli. Loss of AChE activity may lead to a range of effects resulting from excessive nervous stimulation and culminating in respiratory failure and death.

The pesticides employed to kill weeds are called herbicides or weedicides. In early 20 th century various inorganic salts were used to control weeds. But most of the herbicides are organic compounds (Kumar Hemant and Gupta A.B., 1997; Nath Ravindra, Banerajee V., 1999; Grillitsch et al., 1999; Mc. Kim J. M., and Lien G. J., 2001; Hoff et al., 2003; Mc Donald M.D. and Grosell M., 2006).

Glyphosate kills plants by inhibiting the activity of the enzyme 5- enolpyruvylshikimic acid-3-phosphate synthesis (EPSP), which is necessary for the formation of the aromatic amino acids tyrosine, tryptophan, and phenylalanine. These amino acids are important in the synthesis of proteins that link primary and secondary metabolism. Glyphosate also act as a competitive inhibitor of phosphoenolpyruvate (PEP), which is one of the precursors to aromatic amino acid synthesis. It also affects other biochemical processes, and, although these effects are considered secondary, they may be important in the total lethal action of glyphosate (Monsanto Company 1985; Sawada Y., Nagai Y., 1987; Moses M., 1989; Tai T. et al., 1990; Tominack et

al., 1991 Talbot A.R., et al., 1991; Ayoola S.O., 2008).

Choice of test organism

The introduction of pesticides in water can decrease the health of natural aquatic ecosystems. To determine the "health status" of a river or dam, it is important to consider an effective and reliable test organism. The following criteria must be taken into account when choosing a test organism (Rand and Petrocelli 1985; Nussey 1994; Wepener 1997).

- since sensitivities vary among species, species with a broad range of sensitivities should be used whenever possible.
- must be a widely available and abundant species.
- species should be susceptible to routine maintenance in the laboratory and for culturing and rearing them in the laboratory so that chronic toxicity tests can be conducted.
- a test organism must be part of a food chain which can influence people or any other important species.
- in a biological community a test organism must show a low variability on genetic and niche level.
- a test organism must easily be identified because taxonomic indistinctness can influence data interpretation.
- an area's ecological balance cannot be disturbed when the test organism is removed from its natural habitat.

- test organisms must also be available throughout the year and not be too expensive.
- a sufficient number of test organisms of the same size and age must be available for tests.
- a test organism must be tested effortlessly, without using high-priced instruments and must not be labor-intensive.
- a test organism must accumulate pollutants rapidly thereby reflecting environmental levels of the pollutant, this facilitates a better understanding of their distribution.

Aquatic organisms like fishes, integrate all the stresses placed on the aquatic ecosystem. These organisms reflect the combined effects over periods of time. Fishes are a measure of environmental health, because everything that happens on the landscape goes into the rivers. For many years fishes have been valued as excellent indicators of water quality. Fishes have also been the most popular test organism because they are presumed to be the best understood organism in the aquatic environment (Roux D.J., 1994 Stein, J. E., et al., 1992; De La Torre et al., 2000). Because fishes are in direct contact with their surrounding environment, any change in the environment will be reflected as changes in physiological processes and survival (Fernando M.D. and E.A., Moliner 1991; Barton B.A., and G. K., Iwama 1991; Wilson R.W. and Taylor E.W., 1993; Wepener 1997). There are a lot of advantages in using fishes as an experimental animal.

- fish populations and individuals generally remain in the same area.

- fishes represent a wide range of tolerance from very sensitive to highly tolerant.
- the taxonomy of fishes is well established.
- fishes are easy to collect with the right equipment.
- lower cost and shorter developmental time.

Hence, in the present study *Channa punctatus* were chosen as test organism.

Fishery losses by pesticides

Many fish kills have been attributed to pesticides. In the 1960s and early 1970s, 18% of the reported fish kills were attributed to pesticides. Among major fish kills in which more than 100,000 fishes were killed, pesticides were the sixth ranking cause of death (responsible for 3.7% of the deaths). In aquatic systems pesticides cause fishery losses in several ways. These include high pesticide concentration in water, low level doses that may kill highly susceptible fishes, the elimination of essential fish food, or the reduction of dissolved oxygen level in the water due to the decomposition of aquatic animals and plants killed by pesticides. Each year large numbers of fishes are killed by pesticides based on EPA data, we calculated that from 1994-2004 the number of fish kills due to all factors was 145 million fish per year. (EPA 1994 - 2004). Studies have shown that high dosages of herbicides, insecticides, pesticides for weed and pest control are extremely toxic to fish (Corbett J.R., 1974; Jeyaratnam J. et al., 1982; Monsanto Company

1985; Cranmer M.F., 1986; Sawada Y., Nagai 1987; Asztalos et al., 1990; Tominack et al., 1991; Wesseling 1993; O'Malley M., 1997).

Purpose of physiological investigations:-

Fish live in very intimate contact with their environment and are therefore very susceptible to physiochemical changes, which may be reflected in their blood components. In fishes, exposure to chemical pollutants can induced either increase or decrease in haematological levels (Agrawal S.J., Srivastava A.K., 1980; Gill T.S. and Pant J.E., 1981; Hughes 1981; Sharma R.C. and Gupta N., 1984; Kori-Siakpere O., 1985; Pascoe et al., 1986; Kori-Siakpere 1991; Samprath et al., 1993; Annune P.A., et al., 1994a; Shastry K.V., et al., 1997; Neumosok J. G, Hughes G. M., 1998; Das et al., 2004). The count of red blood cell is a stable index and the fish body tries to maintain this count with in the limits for certain physiological standards using various physiological mechanism of compensation. Studies have shown that when the water quality is affected by contaminants, many physiological changes will be reflected in the values of haematological parameters (Haley 1979; Guzelian 1982; Suege J. and Bluzat R., 1983; Matsumura F., 1985; Hoar S.K, et al., 1986; Moses M., 1989; Tai T., et al., 1990; Edwards 1991; Talbot 1991, Mbiapo Youovop 1993). Blood cell responses are important indicators of changes in the internal and / or external environment of animals. The percentage of haemoglobin may also be affected by the toxicants. Any changes in blood parameters of fishes which occur because of the internal injuries can be used to determine the toxicity level of pesticides. (Fry F.E. 1971; Matthiessen P., 1981; Lakota et al., 1983; Golow A.A. and Godzi T. A. 1994; Alabaster J.S. and R., Lloyd 1982; Gaur et al.,

1992; Extoxnet 1995).

Stressors induce some changes that alter the homeostasis of the animals. The stressors first activate the chromaffin cells present in the walls of the cardinal veins and in some cases the heart and kidneys of the teleosts (Kleerekoper et al., 1973; Rand 1977; Mazeaud 1981; Natarajan 1981; Peakall 1992; Weber D.N., Spieler R.E. 1994; Dodson et al., 1995; Beauvais S.B., and Jones S.B. 1999; Beauvais S.L. et al., 2000; Brewer et al., 2001). Chromaffin cells release adrenaline and a small amount of nor adrenaline that stimulates the liver glycogen into blood glucose and the utilization of glucose by muscles. Blood sugar has a direct correlation to metabolism. The organs involved in maintaining the proper blood glucose levels is liver. Any changes caused by toxicants alter the homeostasis as affecting the blood parameters and functions of liver of the fishes (Hochachka 1978; Shastry K.V. and Siddiqui A. A. 1982; Ghosh 1987; Ghosh 1989; Jyothi B. and Natrajan G.M. 1999; Kumar Hemant and Gupta A.B. 1997).

The biggest sources of water in Bundelkhand region is Betwa river. Betwa rises from M.P. and meets the river Yamuna in Hamirpur district in U.P. The Betwa basin includes parts of a number of district in Bundelkhand region, like Sagar, Tikamgarh, Chhatarpur, Lalitpur, Jhansi, Jalaun and Hamirpur. Except Betwa river Bundelkhand have rich source of other water reservoir as Cane river, Matateela dam and Pahunj river. These water reservoirs have become rich source of fish farming and valuable sources of fish food reaching to different areas of our country. The *Clarius batrachus*, *Leabio rohita*, *Heteropneustes fossilis* and *Channa punctatus* are some fishes mainly used for food.

These fishes are continuously challenged or stressed by lethal and sub lethal levels of contaminants. Although several studies have been carried out on physiological parameters exposed to different toxicants but still a little work has been done in this region. Fish haematology is used in recent to detect the stress caused by pesticides. Since glyphosate, phosphamidon, imidacloprid and metasystox are extensively used for pest and weed eradication in Bundelkhand region, it is necessary to study their hazardous effect on the aquatic system especially on fresh water fishes. Due to their food values *Channa punctatus* are in high demand. Thus the present investigations of haematological and biochemical parameters are undertaken to evaluate the effect of the (glyphosate, phosphamidon, imidacloprid and metasystox) toxicants in fresh water teleosts *Channa punctatus*.

Chapter - 2

REVIEW OF LITERATURE

REVIEW OF LITERATURE

The desire of innovation in man has contributed to pollution in the life and ecology of plants, fishes and other animals and microbes. Increased demand for food and other useful material has led to the chemicalization of agriculture and we have reached on such a stage that modern agriculture is dependent on high yielding varieties, which can only be grown under the influence of fertilizers and pesticides. Pesticides are the man made chemicals which are being used to produce enough cheap food. In India, 90,000 millions of technical grade pesticides are used annually to control pests and plant diseases. The steady flow of agricultural effluent discharge into water bodies increases the range of pollutants, which becomes evident when considering toxic pollution (Dhilon S. S., and Gupta A. K., 1983; Forstner U., and Prosi F 1989; Peakall D. 1992; Braunbeck T., 1994; Heath A. G. 1995; Mason C.F. 1996; Heath A. G. 1996; Das R. 2000; Santhakumar et al., 2000; Alam, M. N. and D. N. Sadhu 2001; Mishra et al., 2001; Lata S., et al., 2001; Singh S and D. N. Sadhu. 2001; Alam M. N., 2002; Patnail et al., 2002; Tilik 2005; Radha et al., 2005). Indiscriminate use of pesticides, careless handling, accidental spillage, or discharges of treated effluents into natural waterways have harmful effects on the fish population and other forms of aquatic life and may contribute long term effects in the environment. Water pollution is globally as a potential threat to both human and other animal populations (Verma S.R., et al 1990; Heath AG., 1995; Heath AG., 1996; Braunbeck, T., 1994; Mason C.F., 1996; Hrudey et al., 1996). Pesticides have been the focus of interest for environmentalist because of their wide spread use, transport, distribution in the

ecosystem. Chemicals with potential for wild life damage of aquatic life include heavy metals, organochlorine, organophosphate carbamate and cypermethrin. (Flos et al., 1987; Abidi R. and U.S., Srivastava 1988; Benerjee G., and Rajendranath 1990; Kumar B. and Banerji V., 1990; Kori-Siakpere O., 1991; Mishra B. K., 1993; Akira Kakuno and Jiro Koyama 1994; Bala et al., 1994; Martinez et al., 1994; Ahmad et al., 1995; Gupta et al., 1995; Khattak I. U. D., Hafeez, M. A. 1996; Kumar Hemant and Gupta A.B., 1997; Alkahem et al., 1998; Dethloff et al., 1999 Kori-Siakpere O., Egor V.E. 1999; Dhembare A.J., and Pondha G.M. 2000; Chandra et al., 2001; Aguiwo J.N. 2002; Sherry P. M., and Abidi Z. Z., 2002; Das et al., 2004; Kuruppasamy et al., 2005; Ajani F., 2008; Khalid et al., 2008). Reports of the presence of pesticides in water and food materials have aroused great concern among the scientists and bio researchers, hence led to studies on their toxic potential. There are numerous bio researchers who gave the creative evidence of presence of pesticides in aquatic media (Bhatia, H.L. 1972; Verma, S.R., et al., 1979; Sehgal and M.K. Bhasin 1985; Jhon, P.J. Rathor. And A. Prakash 1989; Verma, S.R., et al 1990; Rajendranath, T. and G. Benerji., 1991; Jain, R. and K.D. Mishra. 1993; Sexena, V., et al., 1997; Das, R. 1998; Santhakumar, M., et al., 2000; Alam, M. N., 2002; Radha, G.S., et al., 2005).

Now a days organophosphorus pesticides are widely used to control the variety of agricultural pests (Khangarot B.S. and P.K. Ray 1987a; Mukherjee M.K. and S.K., Knor. 1984; Mishra U.K. 1988; Ghosh and Chatterji 1989; Khangarot B.S., et al., 1996 Mishra et al., 1998; Tilik K.S., 2005; Trivedi, S. and D. N., Saksena 1999; Benerji G. and Rajendranath 1999; Dhembare A.J. and Pondha, G.M., 2000;

D. S. et al., 2001 Singh S. and D. N., Sadhu. 2001; Delor M.E., et al 2002). Various types of phosphorus group are present in the major class of organophosphorus pesticides. Like as Phosphate group, *O* - alkyl phosphorothioate, phosphorodithioate, *S*-alkyl phosphorothioate, phosphoramidate etc. At least 100 organophosphorus pesticides have been reviewed by WHO for consideration as agents for the control of disease vectors. A large number have been reviewed by the FAO/WHO Joint Meetings on Pesticide Residues. Unlike many compounds scrutinized by the IPCS, these compounds are designed to be toxic for certain pests and are added deliberately to the environment (Gill et al., 1990; Benerjee G., and Rajendranath 1990; Balasubramanian S. and Ramaswami M. 1991; Gaur K Pandey Surendra D 1992; Chandrasekhar S and N., Natrajan 1993; Chaturvedi L.D., and Agrawal K., 1993; Chandrasekhar and S., Jayabalan 1993; Sadhu D.N., 1993; Bhattacharya Lata 1994; Gupta et al., 1995 ; Fryday et al., 1996; Kumar et al., 1997; Kumar Hemant and Gupta A.B., 1997; Jebakumar et al. 1997; Benerji G., and Rajendranath 1999; Nath. Ravindra, Banerajee V. 1999; Armbrust K.L., 2000 Delor et al., 2000; Dhembare A.J. and Pondha G.M. 2000; Kamble G.B. and Muley D.V. 2000; Jhosi P. and Deep H. 2002; Medda et al., 2002; Noori et al., 2003; Ayoola S.O. 2008). Although the organophosphorus pesticides are used to control the insect, pests but they are harmful to other non target organism like fishes as they are continually added to water bodies. Several reports of the excessive use of organophosphorus pesticides and their toxicity on fishes and other aquatic animals are available (Charistoforides C., & Hedley-Whyte, J.1969; Blaxhall P. C.1972; Ranke B., and Rybicke 1975; Lone K..P. and M. Y. Javed; 1976 Ransfold K. D.1978; Agrawal S. J. and A. K.

Srivastava 1980; Pandey et al., 1980; Jagdish et al., 1981; Goel et al., 1982; Grues et al., 1983; Mishra J., Shrivastava A. K. 1983; Natarajan G.M. 1983; Kulshrestha S.K. and Arora A.1984; Jai Nath et al., 1984; Dange A.D., 1986; Khangarot B.S., and P.K. Ray 1988e; Ghosh T. K., 1989; Mahesh et al., 1989; Gill et al.,1990 Benerjee G., and Rajendranath 1990; Balasubramanian S. and Ramaswami M. 1991; Gaur K Pandey Surendra D 1992; Chandrasekhar S and N., Natrajan 1993; Chaturvedi L.D., and Agrawal K., 1993; Gupta 1995; Jain R and K.D. Mishra. 1995; Jeba kumar S.R.D. et al., 1997; Mishra 1997; Sastry N.N., et al., 1997; Khangarot et al., 1996; Poonmani R. and B. Dhanakkodi. 1996; Kumar et al., 1997; Kumar Hemant and Gupta A.B., 1997; Jebakumar et al.. 1997; Imbamani, N. and R., Shrivassan 1998; Neumosojk.G, Hughes G. M., 1998; Benerji G., and Rajendranath 1999; Das R., 2000; Dhembare A.J. and Pondha G.M. 2000; Santhakumar et al., 2000; Alam, M. N. and D. N. Sadhu 2001; Mishra D. S. et al., 2001; Lata S., et al., 2001; Singh S and D. N. Sadhu. 2001; Alam M. N., 2002; Delor et al., 2002; Patnail et al., 2002; Tilik 2005; Radha et al., 2005).

Herbicides are also widely used for the control of water plants and weeds which may impede the flow of water during the summer; when sudden heavy rain can cause flooding (Annune et al., 1994). Harmful herbicides enter surface water with the discharge of agricultural wastes. It is biologically very reactive and hence gives rise to both acute and chronic poisoning toxic effects of pesticides on aquatic organism like fishes. Several workers have investigated the physiological disturbance caused by herbicides exposure on fishes (Brooker M. P. and Edwards R. W. 1975; Monsanto Company 1985;

Hoar et al., 1986; Mitchell et al., 1987; Pulla et al., 1987; Sawada Y, Nagai Y., 1987; Mishra U.K. 1988; Moses M., 1989; Anju Kumari and K., Pandey 1990; Tai T, et al., 1990; Talbot et al., 1991; Tominack, et al., 1991; Neskovic et al., 1996; Mishra 1997; Cox C. 1998; Mishra et al., 1998; USDA 1984). Numerous workers have investigated the disturbance in carbohydrate metabolism caused by various toxicants exposure on fishes (Hochachka, P.W., 1978; Srivastava A. K., 1981; Shrivastva A. K. and Singh N. N., 1981; Shastry K.V. and Siddiqui A. A. 1982; Singh H. H., Srivastava A. K., 1982; Shaikha Y.A. and P.K., Hiradher 1985; Ghosh T. K., 1987; Ghosh T. K. 1989; Shobha et al., 1989; Kumar Hemant and Gupta A.B. 1997; Odiete W.O. 1999; Jyothi B. and Natrajan G. M., 1999).

Aquatic toxicologists traditionally have been interested in determining how much of the toxic chemicals, fishes (or other aquatic creators) can tolerate before they die. This is done by standardizing the tolerable concentrations of pesticides during acute toxicity experiments. One of the purpose of the toxicity studies is to conclude whether a potential toxicant harmful to aquatic life, and if so, to find out the relationship between toxicant and concentration and its consequence on aquatic animals (especially fishes). In this manner various bioresarches have been investigated the acute toxicity effects of pesticides, insecticides and herbicides on aquatic animals (Pandey et al., 1980; Jagdish et al., 1981; Goel et al., 1982; Grues et al., 1983; Mishra J., Shrivastava A. K. 1983; Natarajan G.M. 1983; Kulshrestha S.K. and Arora A. 1984; Jai Nath et al., 1984; Dange A.D., 1986; Khangarot B.S., and P.K. Ray 1988e; Ghosh T. K., 1989; Mahesh et al., 1989; Gill et al., 1990 Benerjee G., and Rajendranath 1990;

Balasubramanian S. and Ramaswami M. 1991; Gaur K Pandey Surendra D 1992; Chandrasekhar S and N., Natrajan 1993; Chaturvedi L.D., and Agrawal K., 1993; Chandrasekhar and S., Jayabalan 1993; Sadhu D.N., 1993; Nath. Ravindra, Banerajee V. 1999; Armbrust K.L., 2000 Delor et al., 2000; Dhembare A.J. and Pondha G.M. 2000; Imbamani N., and R Shrivassan 1998; Kamble G.B. and Muley D.V. 2000; Jhosi P. and Deep H. 2002; Medda et al., 2002; Noori et al., 2003; Ayoola S.O. 2008).

The suitable biomarker tests that cover all the biological activities and functions of the organism and that can make it possible to judge which of the functions were abnormal, when the organism has been exposed to toxicants These are blood parameters which considered pathophysiological biomarkers of the whole body (Silbergeld E.K., 1974; Adams S.M., 1990; Peakall D. 1992; Stein, J. E., et al., 1992; De La Torre et al., 2000; Vander et al., 2003). Several works have been carried out on heamatological and biochemical parameters to determine the effected biological activities on test organism because the blood parameters are the best biomarkers to investigate the hazardous effect of pollution cause by various toxicants (Bhatia H.L.1972; Blaxhall P. C. 1972; Mahajan C. L., and Juneja S. 1979; Pandey et al., 1980; Jagdish Mishra and Anil K. Shrivastava 1981; Sastry K.V and K. Sharma 1981; Goel et al., 1982; Bhaskar B.R., and K.S. Rao 1985; Kori-Siakpere O. 1985; Casillas, E. and L.S Smith 1987; Sastry et al., 1988; Mahesh et al., 1989; Birendra Kumar and Banerjee V. 1991; Kori-Siakpere O. 1991; Sastry K.V. and A. Gupta 1994; Gupta et al., 1995; Khattak I. U. D., Hafeez, M. A. 1996; Kori-Siakpere O, Egor V.E. 1999; Nath. Ravindra, Banerajee V.

1999; Dhembare A.J., and Pondha G.M. 2000; Das B. K. and Mukherjee S.C. 2000; Chandra et al., 2001; Jhosi P. and Deep H. 2002; Sherry P. M., and Abidi Z.Z. 2002 Srivastava R.K. and Srivastava S. 2002; Das et al., 2004; Khalid et al., 2008; Ramesh M., and Saravanan M. 2008; Saufy et al., 2007).

Glyphosate, Phosphamidon, Metasystox and Imidacloprid are four toxicants studied in this toxicological study because they are widely used in all over world and Bundelkhand region, but very scanty informations are available regarding Glyphosate, and Imidacloprid intoxication of fish haematology. Several workers have been carried out in phosphamidon and metasystox intoxicated fishes in India except Bundelkhand region. (Comes et al., 1976a; Rueppel et al., 1977; Roy et al., 1977; Folmar et al., 1979; Mahajan and Juneja 1979; Sullivan, T. P., and D. S. Sullivan. 1979; Hildebrand et al., 1980; Roslycky E. B. 1982; Natrajan G.M., 1983; Morrison et al., 1984; Newton M. et.al. 1984; Monsanto and Company 1981; Monsanto Company, 1985; Hildebrand et al., 1986; Mitchell et al., 1987; Servizi et al., 1987; Jhon P.J., Rathor. And A. Prakash 1989 ; Roy et al., 1989b; Santillo et al., 1989a; Anton et al. 1993; E.P.A. 1993; Liu et al., 1993; Marrs, et al., 1993; Roberts R.O. and S.G. Berk. 1993; Minaxi Das and Shambhu Prasad 1994; Giesy 2000; Feng et al., 1990; Nagata 1996; Neskovic. et.al. 1996; Shekhar P., and I., Christy 1996; Matsuda et al., 1998; Linz et al., 1999; Tomizawa M. and Casida J. E. 2000; Williams et al., 2000; Anand Kumar 2001; Sheets et al., 2001; Josif Jhon P. 2007).

Aannd kumar et al., (2001) reported decreased level of Hb%, TEC and PCV after both acute and chronic exposure to phosphamidon

on *Heteropneustes fossilis*.

Ravindra Nath and V., Vanerjee (1999) reported decreased TEC, Hb% and PCV% but TLC was increased significantly in lethal and sub lethal toxicity of rogar pesticide with in 24, 96 hours and 7 days exposure. Anju kumari and A.pandey (1990) reported the decrease in Hb content TEC and PCV in *Clarius batrachus* intoxication of teficide and butachor follow acute toxicity of 48 hours according to Chandshekhar S. and N., Jayabalan (1993) Hb % and PCV decreased after exposed to sublethal concentration of endosulphan for different period used like as 7, 14 and 21days. The decrease level in Hb %, TEC, PCV and increase in TLC were reported by several workers using oreganophosphorus pesticides and other toxicant in acute toxicity bioassay experiment in fishes (Mukhopadhyay and Dahadrai 1980; Jagdish Mishra and Anil K. Shrivastava 1981; Kori-Siakpere 1985; Chakraporti P. 1986; Homechaudhuri 1986; Anju kumari and A.pandey 1990; Mishra B. K. 1993; Nath. Ravindra and Banerajee V., 1999; Kori et al., 2008; Ajani F. 2008). Similar responses of Hb %, TEC and PCV were also observed by many authors in chronic cases using oreganophosphorus pesticides and other toxicants in fishes (Sastry K.V. and K., Sharma 1982; Natrajan G.M. 1983; Pandey 1984; Shastry et al., 1984; Shastry et al., 1984; Gill T.S. and Pant J.C. 1985; Rajeshwari et al., 1989; Mala and Shreak 1990; Ruparelia et al., 1991; Patel S.K. and P.G. Parmer 1993; Chandshekhar S. and N. Jayabalan 1993; Mughal et al., 1993; Verma G.P. and Pranamita Panighari 1998; Alkaham et al., 1998; Kori-Siakpere et al., 1999; Nath. Ravindra, Banerajee V. 1999; Anand Kumar 2001; Muhammad Atamanalp and Telat Yanik 2002;

Nuri et al., 2003; Atef M. Al. Atter 2005; Vutukuru S.S. 2005; Shah S.L., 2006; Velisek et al 2006 a, b; Ajani F., 2008; Khalid et al., 2008; Ramesh M. and Saravanan M. 2008; Khalid et al., 2008).

The increase level of TLC was also observed by several bioresearchers using various toxicants (Shastry K.V and K. Sharma 1982; Pandey 1984; Shastry et al., 1984; Gill T.S. and Pant J.C. 1985; Kori-Siakpere O. 1985; Chakraparti P. 1986; Homechaudhuri 1986; Bielinska 1987; Rajeshwari et al., 1989; Ruparelia et al., 1991; Thakur G. K. and P. K. Pandey 1991; Chandshekhar S. and N. Jayabalan 1993; Mishra B. K. 1993; Mughal et al., 1993; Patel S.K. and P.G. Parmer 1993; Kori-Siakpere 1999; Nath. Ravindra, Banerajee V. 1999; Anand Kumar 2001; Muhammad Atamanalp and Telat Yanik 2002; Atef M. Al. Atter 2005; Shah S.L. 2006; Ramesh M. and Saravanan M. 2008).

Minaxi das and Shambhu Prasad (1994) reported increase level of TEC (total erythrocytes count) and Hb% (haemoglobin percentage) using metasystox during 48 hours exposure on *Heteropneustes fossilis*. Increased parameters like as TEC and Hb% which originate in this case have also been reported earlier by Mahajan and Juneja 1979; Dhilon and Gupta 1983; Mishra and Shrivastava 1985; Ghosh and Chatterjee; 1989 during acute toxic exposure. G.M. Natrajan (1983) also reported the increase in total RBC count, PCV, and Hb% in *Heteropneustes fossilis* following chronic toxicity with 30 days exposure of metasystox. Similar results of haematological effects on fresh water fishes using different pesticides have also been established prior (Valicre E.J., and Stickney C.J., 1999; Kumar B.K., 1991; Lavin 1992; Kuruppasamy et al., 2005). Few workers are reported the

decrease level of TLC into toxicants on fishes following exposure investigation Muhammad Atamanalp and Telat Yanik 2002 Verma G.P. and Pranamita Panighari 1998; Khalid et al., 2008 Maule A.G., Schreck C.B. 1990.

Malla et al., (2009) reported increased value of ESR after acute exposure of 24, 48, 72, and 96 hours intoxication of chloropyrifos on *Channa punctatus*. Similar reports have also been done by copious bio-researchers using different toxicant in both acute and chronic toxicity (Kumar B., and Banerjee V. 1990; Singh S., and Bhati DPS. 1991; Chaturvedi L.D., and Agrawal K.1993; Bala Sashi et al., 1994; Goel K.A., and Maya 1996; Nath. Ravindra and Banerajee V. 1999). Nuri at al., (2003) reported the MCHC decreased with increase cypermethrin concentrations, but MCV level increased and MCH was not affected with exposure of deferent cypermethrin concentrations (Van Vuren, J.H.J. 1986). The various alterations in values of MCH, MCHC and MCV were also carried out during acute and chronic toxicity bioassay investigated by many workers (Verma et al., 1979; Jagdish Mishra and Anil K. Shrivastava 1981; Natrajan G.M. 1983; Pandey 1984; Gill and Pant 1985; Kori-Siakpere O. 1985; Chakraporti P. 1986; Homechaudhuri 1986; Bielinska 1987 Ramawamy M. and G.T. Reddy 1988; Bradury and Coats 1989; Rajeshwari et al., 1989; Anju kumari and A.pandey 1990; Mala and Shreak 1990; Singh H.S. and Reddy T.V. 1990; Ruparelia et al., 1991; Mishra B. K., 1993; Patel S.K. and P.G. Parmer 1993; Alkaham et al., 1998; Verma G.P. and Pranamita Panighari 1998; Sopinska and Guz 1998; Kori-Siakpere 1999; Nath Ravindra, Banerajee V. 1999; Das and Mukherjee 2000; Muhammad Atamanalp and Telat Yanik 2002; Nuri et al., 2003;

Svobodova et al., 2003; Atef M. Al. Atter 2005; Prashanth et al., 2005; Vutukuru S.S. 2005; Dobsikova et al., 2006; Velisek et al 2006 a;b 2007; Ajani F. 2008; Khalid et al., 2008; Velisek et al., 2009).

Kumar Hemant and A.B., Gupta (1997) reported the increased level of glucose after acute and chronic exposure of glyphosate and metasystox in *Heteropneustes fossilis* (Verma G.P. and Pranamita Panighari 1998; Khalid et al., 2008).

Chandrasekhar S. and N., Natrajan (1993) reported the increase in blood glucose in fishes exposed to endosulfan in *Heteropneustes fossilis*. Other workers also reported hyperglycemic condition in fishes. (Srivastava A.K., 1981; Nemcsok and Bones 1982; Singh H.H., Srivastava A. K. 1982; Ghosh, T. K. ,1989; Nataragan G. M., 1989; Medda et al., 1993; Kumar Hemant and A.B., Gupta 1997; Jyothi B. and G. Narayan 1999; Luskova et al., 2002; Saufy H., et al., 2007).

Temperature of aquatic environment is important for ensuring survival, distribution and normal metabolism of fishes (Forghally et al., 1973). O₂ level, Salinity or other environmental factors also caused disturbance in hormones, metabolic pathways, enzymes and behaviors of fishes (Gubbins et al., 2000). Haematological parameters are also valuable tools for monitoring the health of fishes and they are affected by many endogenous and exogenous factors (Coutant C.C., 1972; Connors et al., 1978; Wilson R.W. and Taylor, E.W. 1993; Howerton R., 2001).

Numerous bioresearchers have been reported the seasonal variations on blood parameters in fishes and other animals (Charistoforides, C. & Hedley-Whyte, J., 1969; Bridges, D. W., et al.,

1976; Cech, J.J. and D.E. Wohlschlag, 1981; Cech, J.J. and D.E. Wohlschlag, 1981; Raizada, M.N., K.K Jain and S. Raizada, 1983; Harding, J and Hogland, L. B., 1984; Joshi, P.C., 1989; Denton, J.E. and M.K., Yousef, 1995; Collazos, M.E., et al., 1998).


Chapter - 3

EXPERIMENTAL DESIGN AND METHODOLOGY

EXPERIMENTAL DESIGN AND METHODOLOGY

Toxicity tests are experiments or trials designed to assess or evaluate the concentration of toxicants and the duration of exposure required to produce a criterion effect. The aquatic toxicity tests are frequently referred to as bioassays. In order to examine the various effects associated with various length of exposure, the ideal and most widely accepted testing frame work includes acute, sub acute and long term toxicity tests. The acute and sub acute toxicity tests are time bound experiments and these tests were carried out in the present investigation to evaluate the response of the toxicants. The term acute can be used to define either the exposure or the response with in short duration exposure. The results of acute toxicity test are represent in terms of LC_{50} (median lethal concentration) for aquatic species. The LC_{50} value represents the ppm or mg/l concentration of the toxicant that would kill 50% of population of a particular aquatic species. The LC_{50} is the usual method of reporting acute toxicity results. The acute toxicity data are important and beneficial in the fixation of sub lethal concentration for chronic toxicity tests.

Sub acute toxicity is also known as chronic toxicity. This study involves a stimulus which is less severe than the acute stimulus and produces a response in longer time and may become chronic. In this the fishes are exposed to chemicals at levels much lower than those that are acutely fatal but they are exposed over a longer period of time. To assess the native of the toxic effects in more realistic situations sub acute toxicity test were conducted.



*All the acute and chronic toxicity experiments were carried out during winter, summer and rainy seasons. Comparisons of different haematological and bio chemical parameters were presented graphically among different season.

Acute toxicity tests

A. Collection of water sample: -

Chlorine free tap water was used through out the course of the experiment. The physiological characters of water sample like the temperature of the test medium, dissolve oxygen, alkinity, hardness and specific conductivity were tested in the Zonal laboratory U.P. Jal Nigam Babina, Jhansi.

B. Collection of fish Sample: -

During the whole toxicological investigation the fishes *Channa punctatus* were used. These are carnivorous fishes feeding upon insects crustaceans and invertebrates. These fishes are supposed to be good, nutritive fish and easily available in living condition through out the year and are good models for experimental work. The fishes were collected from different water bodies of Bundelkhand region with the help of professional fisher man. Live and healthy fishes were used in all the toxicological investigations. The selected fishes were checked against injury, infection or diseases by keeping in 0.2% of potassium permanganate solution for 1-2 minutes. They were kept in glass aquaria having a capacity of more than 40 liters. All over the experiment water was changed daily. The fishes were acclimatized in laboratory condition for 6-10 days. During acclimatization the fishes were fed egg albumin, earth worms and small insects.

C. Preparation of toxicants: -

The four toxicants namely glyphosate, phosphamidon, metasystox, and imidacloprid were selected in the present experiment

which are used in Bundelkhand region. Their trade name, chemical formula, emulsifiable / soluble concentrations and manufacture's name are given in chapter 4. The solution of toxicant was prepared by dissolving the toxicant in 100 ml of distilled water. From this stock solution desired concentrations of each toxicants were added in test medium. To determine the LC_{50} value several concentrations of toxicants were prepared separately. For example for the preparation of 0.025 ml/liter first we dissolve 2.5 ml of toxicant in 100 ml of distilled water. Then we add 1 ml of that solution in one liter of test medium. If the jar contains 5 liter tap water we add 5 ml of stock solution now concentration will be 0.025 ml/liter.

D. determination of LC_{50} value: -

After acclimatization feeding was stopped 24 hrs prior and during the exposure period. For the accurate determination of LC_{50} values two exploratory and one definitive test were conducted. During initial exploratory examination two concentrations were preferred to guess supposed mortality between zero percent to 100 percent. The two concentrations (lower and higher) of toxicants were introduced in two different jars containing five fishes each. In next (second) exploratory or range finding test 4 or 5 concentrations were taken to get narrow range of concentration for LC_{50} determination. Four or five fishes were exposed against each concentration. From the derivation of second range finding examination several (7-9) different concentrations were preferred for definitive test and ten fishes were exposed to each concentration. The mortality was recorded after a period of 24, 48, 72 and 96 hrs and dead fishes were removed when observed. The concentrations from the definitive test were employed

to determine the LC_{50} values by plotting a dose response curve between percent mortality and concentrations of toxicants (Direct interpolation method). A line was drawn between the point represent the % mortality and concentrations. The concentrations at which this line crosses the 50% lethality line was the actual lethal concentration of toxicant.

E. Collection of blood sample for Haematological & Biochemical parameters:-

In order to find out the effect of each toxicant for haematological and biochemical parameters blood was collected from the fishes of treated and control groups. The experimental plane is given below -

For acute toxicity test

After calculating the LC_{50} values 40 healthy fishes from different water bodies were collected, acclimatized and divided into five groups (A, B, C, D, and E) of eight fishes each. The LC_{50} concentrations of toxicants at 96, 72, 48, 24 hours were added to group D, C, B and A respectively. The 5th group maintained in water without toxicants served as control group. After completing the acute toxicity experiments blood was collected in vials, from each group by severing the caudal peduncle. Small quantity of blood was transferred in oxalated vials and remaining blood was centrifuged to separate the serum. Blood parameters were measured by the method described later.

For sub acute toxicity test: -

For chronic toxicity test 60 fishes were collected from market and washed with 0.2% KMnO_4 to avoid any dermal infection and acclimatized at least 10 days in laboratory condition. The fishes were divided into two groups of 30 fishes each. Group one was exposed to sub lethal concentration (1/10 of 96 hours LC_{50}) of toxicants. The second group was kept as untreated control. After 15, 30 and 45 days 10 fishes from each group were sacrificed, blood was collected and serum was separated for the testing of blood parameters.

Methodology**Preparation of oxalated vials:-**

- | | |
|----------------------|--------|
| 1. Ammonium oxalate | 0.8gms |
| 2. Potassium oxalate | 1.2gms |
| 3. Natural formalin | 0.1ml |
| 4. Distilled water | 100ml |

Mix both type of oxalates and add 0.1 ml of natural formalin. After ward transfer this mixture into 100 ml of distilled water. Hence prepared solution was poured in vials and incubate for 10 minutes. Consequently oxalated vials are ready for the use of blood collection.

Estimation of haemoglobin**Apparatus: -**

Haemoglobinometer consisting of two comparator with glass standards, a cylindrical Hb tube marked both gram and percentage figures and Hb pipette marked at 20 cumm, 0.1N HCl and distilled

water.

Principle: -

Hb is converted into acid haematin by HCl. The brown colour of the compound is matched against a brown glass standard in a comparator.

Procedure: -

1. Fill the Hb pipette with non oxalated blood exactly up to 20 cumm mark by sucking. If a slight excess is drawn in it may be removed by touching the point of the pipette with the finger or gauge. Hb pipette or capillary should be filled carefully with out any bubbles.
2. Empty the pipette into graduated tube already filled by HCl up to the bottom graduation line (mark 2). Rinse the pipette at least three times by drawing in and discharging the blood acid mixture.
3. Mix the acid haematin solution in the tube with the glass rod and allow the tube to stand for ten minutes to develop the colour of the acid haematin.
4. Now dilute the solution of acid haematin by adding distilled water drop by drop, stirring the mixture with glass rod. The comparator is held against good day light and addition of water continued till the colour of the solution matches perfectly with that of the standards. Take the reading in grams percent. The bottom of the meniscus is read.

Total erythrocytes count (TEC):-**Apparatus: -**

Red cell pipette, diluting fluid (Hayem's solution), Neubauer's counting chamber with cover slip (haemocytometer), Microscope etc.

Principle: -

Blood is diluted exactly 1:200 with a special pipette using an isotonic diluted fluid (Hayem's solution) for the preservation of the corpuscles. This diluted blood is placed in special RBC counting chamber (Neubauer's chamber). The cells are counted. The number of cells is multiplied by the appropriate factor to obtain the number of erythrocytes in 1 cu mm of undiluted blood.

Preparation of Hayem's diluting solution: -

This solution has following ingredients.

- | | |
|----------------------|--------|
| 1. Sodium chloride | 0.5gms |
| 2. Sodium phosphate | 2.5gms |
| 3. Mercuric chloride | 0.25ml |
| 4. Distilled water | 100ml |

Mix well both type of salts sodium chloride and sodium phosphate and 0.25 ml mercuric chloride and the prepared combination was poured in 100 ml of distilled water.

Procedure:

1. Take the oxalated blood exactly up to the 0.5 mark in a special pipette (RBC pipette). Immediately draw up diluting fluid up to

the mark 101 by holding the pipette horizontally, thus making a dilution of 1:200 in pipette. Now rotating the pipette between the thumb and forefinger.

2. The cover slip is placed over the Neubauer's chamber so as to cover both the ruled platforms evenly.
3. Before charging the counting chambers liberate few drops of the diluted blood by the pipette and put a small drop at the periphery of cover glass which is already in position. Care must be taken so that the suspension does not flow into the moats on either side nor should be bubble remain beneath the cover glass. Allow 2-3 minutes to pass and check the red cells settle down properly.
4. Count the cells in 80 small squares. These 80 small squares comprise 5 medium size squares, each of which is bounded by a triple line. In counting, the cells which touch the left hand side lines or the upper lines of the square are taken to be with in that square and those which touch the lower or right hand side lines are omitted as out side the square.
5. The total volume of 80 small squares is 1/50 cu mm. The dilution is 1 in 200 so the RBC count is

$$\text{Dilution} \times 1/\text{volume} \times \text{No. of cells counted (N)}$$

$$= 200 \times 50 \times N$$

$$= 10000 \times N \text{ cells / cu mm.}$$

Total leucocytes count (TLC):-**Apparatus: -**

White cell pipette, diluting fluid (Truck's solution), Neubauer's counting chamber with cover slip (haemocytometer), Microscope etc.

Principle: -

Blood is diluted exactly 1:20 with a special pipette using an isotonic solution. The diluted fluid truck's was used for preservation of the corpuscles. The diluted blood is placed in a special counting chamber (Neubauer's Counting Chamber) and the cells in a measured volume are counted. This figure is multiplied by the appropriate factor to obtain the number of leucocytes.

Preparation of Truck's solution: -

This solution has following ingredients.

- | | |
|------------------------|--------|
| 1. Glacial acetic acid | 3 ml |
| 2. Gentian violet | 1.0 ml |
| 3. Distilled water | 97ml |

Mix glacial acetic acid in 97 ml of distilled water than add gentian violet.

Procedure:-

1. Draw oxalated blood in a clean dry pipette up to the 0.5 mark. Suck the WBC diluting fluid up to the mark 11. Mix it thoroughly and charge the counting chamber as described earlier. Allow 2-3 minutes to pass and check that all the cells settle down properly.

2. Count the white blood cells in the 4 corners of the chamber. Each of the corners (single ruled square) is divided into 16 smaller squares. Each four large corner squares (16 smaller squares) has an area of 1 sq mm. In counting the cells which touch the left hand side lines or the upper lines of the squares are taken to be with in that square and those which touch the lower and right side are omitted as out side the square.
3. The volume of 4 corner squares is 0.4 cu mm, Dilution 20, Number of cells N then,

$$1\text{cu.mm contain} = \frac{N \times 20}{0.4} = N \times 50$$

Erythrocytes sedimentation rate (ESR):-

Apparatus: -

Wintrobe tube, (Length 110 mm, diameter 3.0 mm, graduation of lower 100 mm. from 0-100) anticoagulant (double oxalate)

Principle: -

If the anticoagulant is added to the blood and the specimen allowed to stand in a tube, red cells slowly sediment to the bottom of the tube leaving clear plasma as the supernatant. The rate of sedimentation estimated under standard conditions is known as the Erythrocytes sedimentation rate.

Procedure:-

The tube is filled up to the 100 mm mark, allowed to stand in a

vertical position at room temperature for one hour and read the fall of the red cells.

Packed cells volume (PCV):-

Apparatus: -

Wintrobe tube, (Length 110 mm, diameter 3.0 mm, graduation of 100 mm. (from 0-100) anticoagulated blood, Centrifuge machine etc.

Principle: -

An oxalated sample of blood is centrifuged to pack red cells to the maximum. The volume of packed cells is determined. The procedure is reliable because its reproducibility.

Procedure:-

1. The wintrobe tube must be clean and dry. Mix the oxalated sample of blood thoroughly for 3 minutes. Fill the blood in the wintrobe tube up to the 100 division marks. There must be no air bubbles.
2. Fill a second Wintrobe tube with either another sample of blood or with water. This tube is to counter balance the first one during centrifugation.
3. Put the tube in the centrifuge and centrifuge it at 3000 r.p.m. for 30 minutes. Take the reading of the packed red cells. The original column of blood in the tube being 100 mm. The volume of packed red cells can be read directly as a percentage.

Mean Corpuscular Haemoglobin (MCH): -

For estimating the average haemoglobin content of a single red cell in microgram following formula was applied.

$$\text{MCH} = \frac{\text{Haemoglobin gm/100ml}}{\text{RBC in millions/Cubic millimeter}} \times 10$$

Mean Corpuscular haemoglobin concentration (MCHC): -

To find out the average haemoglobin percentage following formula can be used.

$$\text{MCHC} = \frac{\text{Haemoglobin in gm/100 ml}}{\text{PCV}} \times 100$$

Mean Corpuscular Volume (MCV): -

The formula for counting the MCV was

$$\text{MCV} = \frac{\text{PCV of /100 ml blood}}{\text{RBC in millions/Cubic millimeter}} \times 10$$

Blood Glucose:-**Apparatus: -**

Autospan diagnostic kit, colorimeter, and distilled water. The kit contains the following reagents.

Reagent 1 (Glucose oxidase, peroxidase, 4-aminoantipyrine, buffer, stabilizers)

Reagent 2 (Diluent Phenol preservative)

Reagent 3 (Dextrose, benzoic acid)

To prepare the working glucose reagent quantitatively, transfer the contents of vial of reagent one and vial of reagent two in a clean black coloured plastic bottle.

Principle: -

Determination of blood sugar is one of the first biochemical estimation to be applied clinically. The abnormalities of blood glucose levels reflect disturbance in physiology.

Glucose is oxidized by glucose oxidase to gluconic acid and hydrogen peroxide. In a subsequent peroxidase catalyzed reaction, the oxygen liberated is accepted by the chromogen system to give a red colored quinoneimine compound. The red colour so developed is measured at 540 nm and is directly proportional to glucose concentration (Silbergeld E.K., 1974; Folin and Wu., 1996).

Procedure: -

We take 3 test tubes and marked it B, S, T respectively. Afterward 20 μ l of serum was taken in test tube marked T and 20 μ l of glucose standard (reagent 3) is added in standard test tube marked S and third tube is kept as blanked marked B. Then add 1.5 ml of working glucose reagent (prepared by mixing reagent 1 and reagent 2

from the kit) into all three test tubes and then incubate them at 37°C for 10 minutes. To stop the reaction add 1.5 ml of distilled water into all three test tubes. Now we measure the absorbance (optical density) of the test and the standard against the blank at 540 nm.

Calculation: -

$$\text{Glucose (mg/dl)} = \frac{\text{Absorbance of test}}{\text{Absorbance of standard}} \times 100$$

Statistical analysis:-

The statistical significance of data was determined by student' t test as heterogeneity of variance was noticed.

Chapter 4

CHEMISTRY OF TOXICANTS

(a.) Chemistry of Toxicants

Glyphosate

Glyphosate is a systemic herbicide that can control most annual and perinnial plants. It controls weeds by inhibiting the synthesis of aromatic amino acids necessary for protein formation in susceptible plants. This compound was first synthesized in 1955 by E. Beriger of CIBA limited Basle, Switzerland. Glyphosate is strongly adsorbed to soil particles, which prevents it from excessive leaching or from being taken-up from the soil by non-target plants. It is degraded primarily by microbial metabolism, but strong adsorption to soil can inhibit microbial metabolism and slow degradation. Photo- and chemical degradation are not significant in the dissipation of Glyphosate from soils. The half-life of Glyphosate ranges from several weeks to years, but averages two months. In water, Glyphosate is rapidly dissipated through adsorption to suspended and bottom sediments, and has a half-life of 12 days to ten weeks. Glyphosate by itself is of relatively higher toxicity to fishes and other aquatic organisms.

Glyphosate, also known by the trade names Roundup for agricultural use, is a broad-spectrum translocated herbicide, used primarily in agricultural applications and for vegetation control in non-crop areas. It is used for aquatic weed control in fish-ponds, lakes, canals, slows running water, etc. (USDA 1984). Formulations of glyphosate include roundup have been extensively investigated for their potential to produce adverse effects in non-target organisms. Glyphosate is soluble in water, and tends to bind tightly to sediment, suspended particulates, organic matter and soil, becoming essentially


unavailable to plants or other aquatic organisms. Since glyphosate developed in the 1970's, there have been much documented cases of adverse effects on fishes and aquatic invertebrates associated its use for the control of aquatic weeds (Giesy et al. 2000).

Glyphosate is perhaps the most important herbicide ever developed. Literature of toxicological and ecotoxicological properties of glyphosate is extremely sparse, considering its importance as herbicide. Generally, glyphosate is slightly toxic to plant, but it may have an impact on the aquatic environment and also on the other aquatic organisms especially on fishes life. Due to this, its toxicity investigation is very important. The study of lethal and sublethal effects is of special importance for toxicological evaluation of compound. It occurs in two isomeric forms α and β in the ratio of 3:7. Its insecticidal property is mostly due to α isomer. It is used as systemic and also as stomach poison.

IUPAC name: N-(phosphonomethyl) glycine

Molecular formula: $C_3H_8NO_5P$

Structural formula:



Its trade names are mostly Roundup^R Armada; Kleenup^R (isopropyl ammonium); Spasor^R; Squadron etc.

Physical Properties: -

Pure Glyphosate is a colourless, odourless, crystalline solid with a melting point of 185 °C and decomposes at 187 °C producing toxic fumes including nitrogen oxides and phosphorus oxides. Solutions of the Glyphosate salts are corrosive to iron or galvanized steel. Pure Glyphosate is slightly soluble in water (12 g/litre at 25 °C), and is practically insoluble in most organic solvents. The alkali-metal and amine salts are readily soluble in water.

Mode of action: -

Glyphosate penetrates the plant leaf cuticle shortly after contact and begins a cell by cell migration to the phloem, from which it is transported throughout the plants. The herbicidal action usually occurs within 7 days and up to 30 for woody plants (Mc Laren and Hart 1995; Monsanto, 1985) Glyphosate's primary herbicidal mode of action is to block the synthesis of aromatic amino acids and the metabolism of phenolic compounds by disrupting the plant's shikimic acid metabolic pathway, leading to the inability of the plant to synthesize protein and produce new plant tissue. This is the only herbicide known to interfere with this particular pathway (McLaren and Hart 1995). A secondary mode of action affects the photosynthetic process, respiration and synthesis of nucleic acids by interacting with a complex series of enzymes which control synthesis of important molecules such as chlorophyll.

Phosphamidon

Phosphamidon is one of the versatile organophosphorus

pesticide extensively used in agricultural operation all over the world since 1956. Phosphamidon was first synthesized in 1955 by E. Benger of CIBA limited, Basle, Switzerland. Technical grade material is dark brown and commercial product is bright violet due to addition of a dye. It occurs in two isomeric forms a-and b-in the ratio of 3:7. Its insecticidal property is mostly due to b-isomer. It is a systemic poison and also acts as stomach poison. It has relatively low contact action. It is commonly used against insects possessing piercing and sucking, and chewing mouth-parts. It is widely used for the control of yellow borer and other sucking pests. There are mainly three routes of entry into water sources. One is from industrial waste or effluent discharged directly into water. A second is by seepage from buried toxic wastes into water supplies. Thirdly, contamination of running water directly or from run-off during spraying operations can occur. Mostly due to rainfall and drainage the pesticide residue reach the water bodies of many non target organism causing toxic effect on fishes.


Primary Use: Insecticide

IUPAC : 2-chloro-2-diethylcarbamoyl-1-methylvinyl dimethyl phosphate

Common name : Phosphamidon

Molecular formula : $C_{10}H_{19}ClNO_5P$

Structural formula :



Trade names : Phosphamidon, Famfos, Dimecron etc.

Physical properties: -

Phosphamidon is a pale yellow to colorless oily liquid with a faint odour. It has a boiling point of 162°C at 1.5 mmHg. Phosphamidon exists as a mixture of 70% cis-isomer and 30% trans-isomer and is corrosive to iron, tinplate and aluminum and it is miscible with water. It is soluble in aromatic hydrocarbons but insoluble in non polar aliphatic hydrocarbons.

Mode of action: -

Phosphamidon poisoning inhibits the activity of superoxide dismutase and increases the lipid peroxidation in several regions in the central nervous system (Brain O' 1987). Delayed neuropathy is initiated by an attack on a nervous tissue system. The target has esterase activity and is called neuropathy target esterase (NTE). The disorder develops not because of loss of esterase activity, but because of a change brought about in the protein molecule that results from the process of ageing of inhibited NTE. The catalytic activity of NTE appears in the nervous tissue, even during the period of development of neuropathy (Edward et al., 1991). Investigations conducted by Extoxnet (1985); Matsumura et al., (1985); Rend G.M. and S.R., Petrocelli (1985) in fishes showed that phosphamidon is potentially neurotoxic because of its ability to inhibit brain NTE activity.

Organophosphorus pesticides exert their acute effects by inhibiting acetyl cholinesterase in the nervous system with subsequent accumulation of toxic levels of acetylcholine. They may also inhibit butyl cholinesterase as well as other esterase. The function of butyl

cholinesterase is unknown, but its inhibition can provide an indication of exposure to an organophosphate.

Metasystox


This pesticide was evaluated in 1965 by Joint Meeting of the FAO Committee on Pesticides in Agriculture and the WHO Expert Committee on Pesticide Residues (FAO/WHO, 1965) under the name of Oxydemeton-methyl. (ODM) Oxydemeton-methyl, S-[2-(ethylsulfinyl)ethyl], which is most commonly sold under the trade name of Metasystox R, is a systemic contact insecticide with approximately 120 tones applied in the states of India during 2004. Approximately 50% of the total was applied to broccoli, and 20% was applied to cauliflower during the year. It was assumed that ODM would be detected in air samples near application sites due to its relatively low vapor pressure (3.80 k Pa), but would not be found long distances from application sites. Dioxydemeton-methyl is a potential transformation product of ODM in air samples.

Primary Use : Insecticide, Oxydemeton-methyl

IUPACname : O,O-dimethyl-S-2-(ethyl-sulfinyl)-ethyl / phosphorothioate

Molecular formula : $C_6H_{15}O_4PS_2$

Structural formula :



Tread name : Metasystox

Physical properties

The molecular formula of metasystox is $C_6H_{15}O_4PS_2$ and its molecular weight is 246.3. it is clear amber-colored liquid (pure compound). The melting point is 10 C (pure compound), boiling point is 106 C/0.01 mm Hg (pure compound) and it is miscible in water.

Mode of action

Oxydemeton-methyl (Metasystox R) is an organophosphate pesticide. The target organ is nervous system and inhibiting cholinesterase enzyme. It binds to the enzyme that is normally responsible for breaking down Ach. When an insect has been poisoned by a cholinesterase inhibitor, the cholinesterase is not available to help breaking down the ACh, and the neurotransmitters continue to cause the neuron to "fire" or send its electric charge. This causes over stimulation of the nervous system, and the insect die.

Imidacloprid

Imidacloprid was discovered in 1984 at Nihon Bayer Agrochem in Japan by screening novel synthetic compounds for a high affinity to the insect nicotinic AChRs receptors, but with low toxicity to vertebrate species (Kagabu 1997). Its molecule includes the insecticidal N-(3pyridinyl) methyl group of nicotine and a nitroimine moiety, Because of their structural similarity to nicotine, imidacloprid and related insecticides (acetamiprid, thiacloprid, thiamethoxam and nitenpyram) were termed neonicotinoids (Tomizawa and Yamamoto, 1993). Imidacloprid was first registered in U.S. in 1994 as a pesticide


and developed for commercial use. Its major manufacturer is Bayer Corporation that markets imidacloprid products with the brand names Admire, Advantage, Confidor, Gaucho, Supreme etc.

Imidacloprid is a systemic chloronicotinyl insecticide that enters the target pest via ingestion or direct contact. In the environment, the principal routes of dissipation for imidacloprid are aqueous photolysis, microbial degradation and uptake by plants. The major degradation product of imidacloprid in the environment is desnitro-imidacloprid. Imidacloprid is used for controlling sucking insects, soil insects, termites, and some chewing insects.

Primary Use : Insecticide

IUPAC name : 1-(6-chloro-3-pyridylmethyl)-N-nitroimidazolidin-2-ylideneamine) is a synthetic active ingredient used in various insecticide

Structure formula



Physical properties:-

It is a colourless crystalline solid, has the molecular formula

$C_9H_{10}ClN_5O_2$ and a molecular weight of 255.7 (Tomlin 2000). Imidacloprid is soluble in water, relatively non-volatile.

Mode of action: -

Imidacloprid, and other insecticides in the nicotinoid chemical family, are "similar to and modeled after the natural nicotine [a tobacco toxin]. Because of their molecular shape, size, and charge, nicotine and nicotinoids fit into receptor molecules in the nervous system that normally receive the molecule acetylcholine. Acetylcholine carries nerve impulses from one nerve cell to another or from a nerve cell to the tissue. Imidacloprid and other nicotinoids irreversibly block acetylcholine receptors. It is used to treat seeds, soil, crops and structures, and is a flea control treatment on domestic pets (Meister, 2000). The toxicity of imidacloprid is based on interference of the neurotransmission in the nicotinic cholinergic nervous system. Imidacloprid binds to the nicotinic acetylcholine receptor (nAChR) at the neuronal and neuromuscular junctions in insects.

(b.) Physical properties of water: -

The chlorine free tap water was used through out the course of the experiment. The physiological characters of water sample like as the temperature of the test medium, pH, dissolve oxygen, alkinity, hardness and specific conductivity were tested in the Zonal laboratory U.P. Jal nigam Babina. Analyses of water have done during all three seasons.

Table : Physical properties of water

S. No.	Water Parameters	Rainy	Summer	Winter
1	Dissolve oxygen mg/liter	7.6	6.2	8.9
2	Alkinity as CaCo ₃ to methyl orange	326	320	308
3	Hardness as Ca	120	111.7	128.7
4	Specific conductivity micro mho	792	765	782
5	Water temprature °C	27	42	14
6	pH value	7.4	6.8	7.2

Chapter - 5

**EFFECT OF GLYPHOSATE
ON BLOOD PARAMETERS OF
*CHANNA PUNCTATUS***

EFFECT OF GLYPHOSATE ON BLOOD PARAMETERS OF *CHANNA PUNCTATUS*

Glyphosate 41% S.L. (trade name Round-up) was used in this study. The chemical composition of this compound is isopropyl amine salt of glyphosate. It is manufactured by Monsanto chemical of India limited, Mumbai.

Results

(a) Acute toxicity bioassay

In first exploratory test 100% mortality was observed in 0.030 ml/l of Glyphosate while no mortality occurred in 0.001 ml/l (Table.1). In second exploratory test four concentrations (0.007, 0.013, 0.019 and 0.025 ml/liter) were taken and mortality was recorded as shown in table 2. Eight different concentrations between 0.007 & 0.025 ml/l were left out for definitive test and mortality data were recorded (Table 3). Then a curve was plotted between percent mortality and concentrations of glyphosate used in definitive test. LC_{50} values were determined by drawing a line intersecting through the concentration of Glyphosate at 50% mortality level (Fig 1 and 2). The values of LC_{50} were estimated as:-

- (1.) 24 hours-0.018 ml/liter
- (2.) 48 hours-0.015 ml/liter
- (3.) 72 hours-0.012 ml/liter
- (4.) 96 hours-0.009 ml/liter

The behavioral changes were observed following exposure to

glyphosate. They show uncoordinated behaviour, agitated or erratic swimming excessive secretion of mucus. The colour of the tested fishes appeared little pale as compared to the normal fishes. Initially exposed fishes came to surface to engulf air frequently. After that the exposed fishes became very weak and settled at the bottom and died.

(b) Haematological & biochemical study:-

The haematological and biochemical parameters of control and glyphosate exposed fishes at LC_{50} concentrations are tabulated (Table 4). The Haemoglobin percentage and total erythrocytes counts were decreased significantly ($P < 0.001$). The Hb% in control fishes was 14.1 gm % where as in treated fishes it was 10.5, 10.7, 10.9 and 11.2 gm % at 24, 48, 72 and 96 hours respectively. The Hb % was lower at 24 hours LC_{50} . Although TEC was less from control fishes but gradually increased as exposure period increased but the difference between treated and control fishes at 96 hours LC_{50} was not significant. A significant decrease was also observed in PCV, MCH & MCV levels ($P < 0.01$). PCV was lowest at 24 hours and highest at 96 hours. MCH levels were also slightly increased from 24 – 96 hours exposure period. At 48 hours LC_{50} MCV was highest but at 96 hours it was lowest following exposure of glyphosate, but all these values were decreased when compared to control fishes. The TLC and MCHC levels increased significantly at $P < 0.01$. ESR was also increased from the control fishes. The ESR was 2.3 mm in control fishes and in herbicides treated fishes it was 3.3, 3.8, 3.8 and 3.6 at 24, 48, 72 and 96 hours LC_{50} , but the difference was insignificant at 48, 72 and 96 hours exposure period.

The table 4 shows that the level of glucose increased significantly ($P < 0.01$) in treated fishes. The level of glucose was 64.10 in control fishes where as fishes exposed to acute toxicity concentrations of Glyphosate showed 72.05, 74.19 74.10, 74.90 mg /100 ml blood of glucose concentration.

In chronic toxicity bioassay (Table 5) the Hb % TEC, PCV, MCH & MCHC decreased significantly ($P < 0.01$) after 15 days, 30 days and 45 days exposure of glyphosate (1/10 of 96 hours LC_{50} concentration). The Hb% was lowest after 15 days exposure period. The levels of other parameters such as TEC, PCV, MCH and MCV were also lowest after 15 days of exposure but gradually increased with increasing exposure period. A significant increase in the level of total leucocytes counts was observed ($P < 0.01$). Although an increased level of ESR was observed when compared with the unexposed fishes but statistically the difference was not significant. The table 5 indicates the changes in blood glucose level of *Channa punctatus*. Glucose level increased significantly ($P < 0.01$) when the fishes were subjected of glyphosate treatment. It was lower after 30 days exposure period.

(c) Study of seasonal variation:-

Seasonal changes of some haematological and biochemical parameters (glucose levels) are presented in the figures 3-11 during acute toxicity bioassay and in figures 12-20 in chronic toxicity bioassay. In acute and chronic toxicity tests the activity of haematological parameters (Hb%, TEC, TLC, ESR, PCV, MCV, MCHC and MCH) levels in blood were lower in late summer season while highest in rainy season. Although the blood parameters like

Hb%, TEC, PCV, MCV, MCH were decreased and TLC, ESR & MCHC were increased from the control fishes as shown in table 04 and 05 but the activity levels of these parameters were low in summer. The similar results have seen as in case of blood glucose level which increased in treated group when compared to untreated fishes but the level in treated and control group was lowest in summer season.

Discussion

(a) Acute toxicity:-

Formulations of Glyphosate including Rodco, Round up and Aquamaster have been extensively investigated (Monsanto Company 1985; Sawada Y., Nagai Y., 1987; Moses M., 1989; Tai et al., 1990; Tominack et al., 1991; Talbot et al., 1991). Glyphosate is used as non selective herbicides for aquatic weed control in ponds lakes and canals. The recommendation of field application rate usually ranges from 1500 and 2000 ppm in Asian countries. According to Wannee Jiraungkoorskul et al., (2002) the LC₅₀ values of glyphosate (48 %) were 17.5, 17.1, 16.9 and 16.8 ppm at 24, 48, 72 and 96 hours after exposure on fresh water fishes *Oreochromis niloticus*. The acute toxicity of glyphosate (62 %) has been investigated by Mitchell et al., (1987) in rainbow trout using different dilutions and physiological conditions of water and the result shows that 96 hours. LC₅₀ value ranges from 15-26 mg /liter. This result was also supported by Ayoola S.O., (2008). He calculated the LC₅₀ values in *Oreochromis niloticus*, at 96 hours which was 1.05 mg/liter. Several works have also been carried out using glyphosate and other herbicides in fresh water fishes

(Brooker, M. P. and Edwards, R. W., 1975; Monsanto Company 1985; Sawada et al., 1987; Moses 1989; Anju Kumari and K., Pandey 1990; Tai et al., 1990).

Kumar Hemant and Gupta (1997) showed the effect of Glyphosate (Isopropyl amine salt of glyphosate) in *Heteropneustes* and estimated 146, 134, 129, & 124 ppm LC₅₀ values at 24, 48, 72 and 96 hours. In the present study the LC₅₀ values were 0.018, 0.015, 0.012 and 0.009 ml/liter at 24, 48, 72 & 96 hours which correlates the LC₅₀ values of other workers. A review of literature revealed that, this is the first study made to evaluate the LC₅₀ values of glyphosate for *Channa punctatus* in Bundelkhand region.

Several abnormal behaviour such as jumping and gulping of air, restlessness, sudden quick movement at the bottom in *Channa punctatus* were similar to the result of Omoniyi et al., (2002); Rehman et al., (2002); and Aguigwo J.N., (2002). The erratic and stressful behaviour of fishes in the present study indicated respiratory impairment, which may be due to the effect of glyphosate herbicides on the gills. (Vivekanandan E., Pandian T.J., 1977; Black et al., 1980; Baldwin et al., 1994; Allin, C.J., and R.W., Wilson. 2000; Weis, J.S., et al., 2001). Light microscopic study of the gills of *Nile tilapia* exposed to glyphosate showed several pathological changes and their frequencies increased with increasing the exposure time (Ayoola S.A., 2008). The fishes came to surface to engulf air. This may be due to low rate of Oxygen uptake under toxicity stress (Wilson, R.W., Taylor, E.W., 1993; Stein, J. E., et al., 1992; Vander et al., 2003).

(b) Haematological and biochemical study:-

Synthetic herbicides are commonly used by farmers to control weeds and nuisance aquatic vegetations around rivers, lakes and reservoirs. However, these herbicides find their way to water bodies, inducing adverse impact on fishes living their in (Tsuda et al., 1997). Since blood forms a unique component between the external and internal environment, toxicant that cause stress in fishes can alter the composition of blood. So that the use of haematological parameters for toxicological research is important for environmental monitoring and fish health conditions. Many works have been conducted on haematological changes of herbicides in fishes (USDA., 1984; Mitchell D.G., et al., 1987; Servizi J.A., et al., 1987; Singh H.S., Reddy T.V., 1990; Neskovic N., et al., 1993; Annune P.A., et al., 1994; Risbourg, S.B., 1995; Rahman M.Z., 2002; Ayoola S.O., 2008). The significance reduction in the values of haemoglobin percentage and TEC after exposure to lethal and sub lethal concentration is an indication of anaemia caused by glyphosate on the exposed fishes. The inhibition in the level of Hb% and TEC were also observed by Kori Siakpers., (2007) using the herbicides paraquet. Significant decrease occurred in the Hb% level may be due to the impairment of oxygen supply to various tissues especially gills. (Ayoola S. 2008; Neskoric N.K., et al., 1996; Rojik J., et al., 1983). Reduced TEC may be as a result of anaemia which is due to haemodilution resulting from impaired osmoregulation across the gill epithelium as reported by Svododa et al., (2001).

Increased in TLC has been attributed to several factors such as increase in thrombocytes and lymphocytes (Agrawal and Shrivastava 1980; Thakur G.K. and P.K., Pandey 1990). Leucocytes are involved

in the regulation of immunological functions of the body (Santa Kumar et al., 2000). An increased in TLCs thus occurs as a protective response to stress (Das B.K., 1998). In the present study PCV value decreased when the fishes are poisoned by toxicant (Gill and Pant 1985). The reduction indicates that the fishes suffer from anaemia or haemodilution (Wedemeyer et al., 1976). Fall in MCV, MCH and increase MCHC values were observed in the present study in acute and chronic conditions. This alteration confirms anaemia in fishes. These alterations also may be due to histopathological changes occur in RBC following exposure to toxicants in *Channa punctatus* and other fresh water fishes (Dalela R.C., et al., 1979; Sastry K.V., and K., Sharma 1981; Dange A.D., 1986; Gill T.S., Pant, J.C., and Pant J., 1988; Hinton D.E., & Lauren D.J., 1990; Birendra Kumar and Banarjee V., 1991; Nowak B., 1992; Poleksic V., et al., 1994; Brusle J., et al., 1996; Ecobichon et al., 1996; Kashiwada, S., et al., 2002; Choudhary et al., 2003; Vander Oost R., et al., 2003; Capkin E., et al., 2006). Increased level in values of ESR may be due to increase concentration of fibrinogen which develops fibrinogenemia due to various pesticides exposure (Singh S. and Bhati DPS 1991; Mala et al., 2009).

Biochemical characteristics are among the important indices of the status of internal environment of the fishes (Nemcsok and Bones 1982; Edsall C.C., 1999). Changes in the biochemical parameters cause changes in metabolism due to the effect of various pollutants. Changes in the activity of several enzymes and carbohydrate metabolism have been observed by several workers (Hochachka, P.W., 1978; Ransfold K. D. 1978; Jagdish et al., 1981; Shrivastva A.

K., and Singh N. N. 1981; Shobha et al., 1989; Chandrasekhar and Jayabalan S. 1993; Jyothi B., and Natrajan G. M. 1999; Prakash et al., 2007). A significant increase in blood glucose level or hyperglycemia occurs in glyphosate treated fishes. The blood glucose level in Indian cat fishes *Heteropneustes fossilis* exposed to sub lethal concentrations of other toxicants were also found to increase considerably (Love R. M., 1970; Dragomirescu et al., 1975; Ransford K. D., 1978; Jagdish et al., 1981; Shrivastva A. K., and Singh N. N. 1981; Shobha et al., 1989; Chandrasekhar and Jayabalan S. 1993; Hemant and Gupta A.B. 1997; Kumar; Prakash et al., 2007).

Increased in blood glucose level may indicate a greater energy requirement of the fishes due to glyphosate stress. The stress may also be viewed as a response of respiratory insufficiency (Rahman M.Z., 2002; Ayoola S.O., 2008). Altered carbohydrate metabolism and impairment in the liver of fishes may be another reason for hyperglycemic condition found in *Channa punctatus* following exposure to glyphosate.

(c) Seasonal variation:-

The data presented in figure 3-11 and 12-20 showed that haemoglobin percentage, TEC, PCV, MCH and MCV decreased but TLC, ESR, MCHC and blood glucose level were increased following exposure to glyphosate from control fishes both in acute and chronic experiment. Although these changes were similar as shown in table 4 and 5, but the concentrations of above parameters in control and treated fishes were lowest in summer and then increase in rainy season. This clearly shows that a process of haemoconcentration

occurs through the late spring and summer and that is followed by haemodilution in the late summer. ESR which is the measure of the concentration of the blood rate, shows down in summer and increases as the blood becomes dilute. A similar response of haemoconcentration and haemodilution is described in fishes and other cold blooded animals like frogs (Schaefer A.A., 1925; Kaplan and Carousa 1956). The lower values in summer may also be due to low rate of oxygen during summer season. (Harding J., and Hogland, L. B., 1984; Joshi P.C., 1989; Akira Kakuno and Jiro Koyama 1994; Dento, J.E., and Yousef M.K., 1995; Gross et al., 1996; Collazos M.E., et al., 1998; Best et al, 2001). The chemical investigations of water in the present study (chapter 4) shows that the low rate or dissolve oxygen in summer season.

Determination of LC₅₀ for Glyphosate on *Channa punctatus*

Table 01: First exploratory test

S.No.	Conc. ml/liter	No. of Fishes	24 Hours		48 Hours		72 Hours		96 Hours	
			M	%M	M	%M	M	%M	M	%M
1.	0.001	5	0	0	0	0	0	0	0	0
2.	0.030	5	5	100	-	-	-	-	-	-

Table 02: Second exploratory test

S.No.	Conc. ml/liter	No. of Fishes	24 Hours		48 Hours		72 Hours		96 Hours	
			M	%M	M	%M	M	%M	M	%M
1.	0.007	5	0	0	0	0	1	20	1	40
2.	0.013	5	1	20	1	40	1	60	1	80
3.	0.019	5	3	60	1	80	1	100	-	-
4.	0.025	5	5	100	-	-	-	-	-	-

Table 03: Definitive test

S.No.	Conc. ml/liter	No. of fishes	24 Hours		48 Hours		72 Hours		96 Hours	
			M	%M	M	%M	M	%M	M	%M
1.	0.008	10	0	0	1	10	1	20	2	40
2.	0.010	10	0	0	2	20	2	40	2	60
3.	0.012	10	1	10	2	30	2	50	2	70
4.	0.014	10	2	20	2	40	2	60	3	90
5.	0.016	10	3	30	3	60	3	90	1	100
6.	0.018	10	5	50	3	80	1	90	1	100
7.	0.020	10	7	70	2	90	1	100	-	-
8.	0.022	10	8	80	2	100	-	-	-	-



Fig 01: LC₅₀ of Glyphosate after 24 hours and 48 hoursFig 02: LC₅₀ of Glyphosate after 72 hours and 96 hours

Table No. 04

Effect of acute toxicity of Glyphosate on selected blood parameters in
fresh water fish *Channa punctatus*

S No.	Parameters	Control	Exposure Period			
			24 Hours	48 Hours	72 Hours	96 Hours
01.	Hb (g/100ml)	14.1 ±0.72	10.7** ±0.52	10.7** ±0.43	10.9** ±0.70	11.2** ±0.25
02.	TECx10 ⁶ /mm ³	3.84 ±0.06	3.46** ±0.05	3.47** ±0.07	3.48** ±0.09	3.45 ±0.02
03.	TLCx10 ³ /mm ³	3.2 ±0.51	3.7* ±0.72	4.2* ±0.3	4.0* ±0.46	3.9* ±0.64
04.	ESR (mm)	2.3 ±0.76	3.3* ±0.28	3.8** ±0.28	3.8** ±0.28	3.6** ±0.28
05.	PCV%	33.0 ±3.0	22.66* ±1.15	24.33* ±1.52	24.66* ±2.88	26.00* ±2.64
06.	MCH pg	36.78 ±2.21	30.85* ±1.21	30.80* ±1.27	31.35* ±1.90	33.26* ±0.67
07.	MCHC %	42.59 ±1.79	47.20* ±0.46	44.13* ±4.21	44.68* ±5.58	43.63* ±4.70
08.	MCV um ³	85.87 ±7.86	70.09* ±10.65	73.80* ±9.73	66.92* ±0.48	65.60* ±9.63
09.	Glucose (Units)	64.10 ±6.82	72.05* ±7.58	74.19* ±8.63	74.10* ±8.49	74.90* ±8.36

* - Significant at P < 0.01; ** - Significant at P < 0.001

Table No. 05
Effect of chronic toxicity of Glyphosate on selected blood parameters
in fresh water fish *Channa punctatus*

S No.	Parameters	15 Days		30 Days		45 Days	
		C	T	C	T	C	T
01.	Hb (g/100ml)	13.9 ±0.60	12.5* ±0.55	14.0 ±0.51	13.4* ±0.50	14.4 ±0.60	13.3* ±0.51
02.	TEC×10 ⁶ /mm ³	3.77 ±0.11	3.58* ±0.91	3.90 ±0.04	3.63* ±0.09	3.98 ±0.09	3.88* ±0.09
03.	TLC×10 ³ /mm ³	3.1 ±0.40	3.8* ±0.37	3.2 ±0.35	4.0* ±0.35	3.5 ±0.40	4.4* ±0.10
04.	ESR (mm)	2.3 ±0.28	3.9** ±0.11	2.5 ±0.11	4.1** ±0.41	2.2 ±0.32	4.2** ±0.23
05.	PCV%	27.33 ±2.30	19.33* ±3.05	30.00 ±3.60	21.66* ±2.51	32.33 ±2.30	24.33* ±2.08
06.	MCH pg	36.85 ±0.47	34.71* ±0.70	36.10 ±1.52	35.43* ±0.67	36.84 ±0.60	36.30* ±0.36
07.	MCHC %	51.08 ±4.71	65.14* ±8.12	47.46 ±6.20	59.73* ±4.55	44.35 ±3.30	56.24* ±3.05
08.	MCV um ³	72.53 ±6.27	53.87* ±7.21	76.93 ±10.17	59.59* ±5.45	83.36 ±6.42	65.87* ±4.05
09.	Glucose (Units)	60.54 ±7.22	72.36* ±10.32	63.34 ±6.71	68.98* ±5.04	65.14 ±5.25	76.75* ±6.47

* - Significant at P < 0.01; ** - Significant at P < 0.001

Graphical comparison of seasonal variation in acute toxicity experiment




Figure 03. Haemoglobin




Figure 04. TEC




Figure 05. TLC




Figure 06. ESR




Figure 07. PCV




Figure 08. MCH




Figure 09. MCHC




Figure 10. MCV




Figure 11. Blood Glucose

Graphical comparison of seasonal variation in chronic toxicity experiment




Figure 12. Haemoglobin




Figure 13. TEC




Figure 14. TLC




Figure 15. ESR




Figure 16. PCV




Figure 17. MCH




Figure 18. MCHC




Figure 19. MCV




Figure 20. Blood glucose

Chapter - 6

**EFFECT OF PHOSPHAMIDON
ON BLOOD PARAMETERS OF
*CHANNA PUNCTATUS***

EFFECT OF PHOSPHAMIDON ON BLOOD PARAMETERS OF *CHANNA PUNCTATUS*

Phosphamidon 41% S.L. (trade name Don 400) insecticide was used in this investigation. The chemical composition of this compound is O-(2-chloro-2-diethylcarbamoyl-1-methyl (vinyl) dimethyl phosphate. It is manufactured by Devi dayal (sale) limited, Reay road, Mumbai.

Results

(a.) Acute toxicity bioassay:-

In first exploratory test 100% mortality was observed in 0.04 ml/liter and no mortality in 0.002 ml/l (Table 06). In second exploratory test mortality due to different concentrations were observed (Table 07). The concentrations were (0.007, 0.014, 0.021 and 0.028 ml/liter). The concentrations between 0.007 ml/l & 0.028 ml/l were left out for definitive test because 100% mortality was observed in 0.028 ml/liter at 48 hours and 20% mortality was seen in 0.007ml/liter at 96 hours. Nine concentrations (0.008, 0.010, 0.012, 0.014, 0.016, 0.018, 0.020, 0.022, 0.024 ml/liter) were selected for definitive test and mortality data were recorded in table 08. Then a curve was plotted between percent mortality and concentrations of phosphamidon using definitive test, LC_{50} values were determined by sketching a line intersecting through the concentration of phosphamidon at 50% mortality altitude. (Fig 21 and 22). The values of LC_{50} were estimated as

(1.) 24 hours-0.023 ml/liter

(2.) 48 hours-0.019 ml/liter

(3.) 72 hours-0.015 ml/liter

(4.) 96 hours-0.011 ml/liter

The behavioural changes were observed following exposure to phosphamidon. It was restlessness, erratic and jerky movement, and loss of balance, jumping and breathing rapidly. Other changes like as the extra secretion and deposition of mucus over the surface of body and fading of body colour has also been observed. The exposed fishes became very weak, settled at the bottom and ultimately died.

(b) Haematological and biochemical study:-

The haematological and biochemical parameters of control (untreated) and phosphamidon exposed fishes at LC_{50} concentrations are tabulated (Table 09). The table 9 shows that the haemoglobin percentage (Hb %) and total erythrocytes counts (TEC) were decreased significantly ($P < 0.001$). Although the values of Hb% and TEC were all the times lesser than untreated fishes but at 72 hours the difference was statistically insignificant. The maximum value of Hb % was found in 24 hours and minimum in 96 hours when compared to control. The table 09 also confirms that the rate of TLC was always greater than untreated fishes, but progressively increased during 24 to 96 hours. MCHC % of blood significantly increased ($P < 0.01$) in fishes after acute exposure to insecticides compared to the corresponding control but it was observed that level of MCHC decreased progressively. The values of ESR were insignificantly increased, but at 96 hours it was significant ($P < 0.01$). The data illustrate that the value of PCV, MCH and MCV were significantly

decreased ($P < 0.01$), but it was also become apparent that the values of PCV, and MCV were lowest in 24 hours exposure and highest in 96 hours intoxication of phosphamidon.

The effect of biochemical parameter (B.glucose) after acute exposure of phosphamidon has been shown in table 09. It was clear that the values of glucose were increased significantly ($P < 0.01$) during treatment of fishes with phosphamidon. The level of blood glucose was 64.7 units in control fishes while 70.07, 73.71, 73.72 and 74.91 units for 24, 48, 72 and 96 hours respectively in treated fishes.

In chronic toxicity bioassay (Table 10) the Hb % TEC, PCV, MCH & MCV were decreased significantly ($P < 0.01$) at 15 days, 30 days and 45 days after exposure at 1/10 of 96 hours of phosphamidon LC_{50} concentration. It was observed that the levels of Hb%, TEC, MCH, PCV and MCV were less than control fishes, but gradually increased with rising exposure periods. The table 10 demonstrates that the levels of TLC, ESR and MCHC were increased significantly at $P < 0.001$.

The level of blood glucose after chronic exposure of phosphamidon has given in table 10. The level of blood glucose was increased significantly at $P < 0.01$. It was lower after 15 days but higher after 45 days exposure period.

(c) Study of seasonal variations:-

The seasonal alterations of selected haematological and biochemical parameters (glucose levels) were obtained and arrange in figure 23-31 during acute toxicity and figure 32-40 during chronic

toxicity bioassay. If comparison was made in control fishes during different seasons, it was found that in acute toxicity cases the activity of haematological parameters (Hb%, TEC, TLC, ESR, PCV, MCV, MCHC, MCH and blood glucose) level in blood were lower in summer season and higher in rainy season. Although the blood parameters like Hb%, TEC, PCV, MCV, MCH were decreased and TLC, ESR, MCHC and blood glucose were increased from the control fishes as shown in table no. 09 & 10, but the activity levels were moderate in winter, slightly high in rainy season and low in summer. Approximately comparable responses were originated in chronic toxicity test.

Discussion

(a) Acute toxicity bioassay:-

Organophosphorus insecticides (phosphamidon) are now extensively used in plant protection operation on accounts of their less persistence in the environment. Various bio researchers have reported the LC_{50} value in different fishes intoxication of phosphamidon depending on their test conditions. Jaya Natha rao et al., (1984) notified the LC_{50} value (16.0 mg/liter) of phosphamidon at 48 hours in *Sarotherodon massambicus*. Sreenivasulu et al., (1991) calculated the 13.0 ppm LC_{50} values of phosphamidon at 96 hours in *Tilapia mossambica*. Sexsena et al., (1997) reported the LC_{50} values 5.54 ppm for 96 hours in *Channa orientalis*, while Jeba kumar (1997) estimated the LC_{50} value 8.0 ppm for 96 hours in *Heteropneustes fossilis*. According to Imbamani and Shrivassan (1998) the LC_{50} value was 2.378 ppm for 96 hours in *S. mossambica*. However Anand Kumar A.,

and Tripathy N.K., (2001) reported 550 ppm and 280 ppm LC₅₀ values for 24 and 96 hours in *Heteropneustes fossilis*. Some other workers have described toxicity of phosphamidon and other insecticides on fresh water fishes. (Mahesh et al., 1989; Sikoki F. D., & Enajekpo H.O.S., 1991; Ghosh R., Shrotri R.V., 1992; Stein et al., 1992; Singh et al., 1992; Canli M., 1996; Luskova V., 1997; Dhembare A.J., and Pondha G.M., 2000; Das B. K. and Mukherjee S. C., 2000; Bhatia, N.P. Sandhu. G.S., Johal M.S., 2002). In present study the LC₅₀ values were 0.011, 0.015, 0.019 and 0.025 ml/liter at 24, 48, 72 and 96 hours. A review of literature revealed that the LC₅₀ values of phosphamidon for *Channa punctatus* in Bundelkhand region was comparable to the estimated LC₅₀ values of previous workers.

The abnormal changes in the behaviour of *Channa punctatus* may be due to the direct manifestation of the disturbances in the physiological mechanism (Fernando, M.D. and E.A. Moliner 1991; Barto B.A., and G. K., Iwama 1991; Wilson, R.W., Taylor E.W., 1993). Swimming movement, restlessness condition, loss of balance, jumping etc were observed in this study may be due to the effect of phosphamidon on central nervous system (CNS) (Brain 1967; Jai Nath and Rao 1984). The rapid breathing was observed when the fishes introduced in toxic environment of phosphamidon. This may be due to hyper excitability which involves considerable energy expenditure and there by making greater demands of oxygen. (Verma et al., 1979; Charistoforides C., & Hedley-Whyte J., 1969; Mishra C.K., Hakim A, Kumar J., 1994 ; Odiete W.O., 1999; Das, P. C., et al., 2003)

(b) Haematological and biochemical study:-

Phosphamidon is one of the exceptional organophosphorus pesticides comprehensively used in agricultural programme in all over the Asian countries. Indiscriminate uses of pesticides result in environmental pollution leading to accidental poisoning of different organism especially fishes in the ecosystem. Most of the chemicals are continuously added to the water bodies through run-off water from agricultural field, causing poisoning of fishes. The physiological stress caused by these chemicals can be studied by haematological parameters, as the actual health condition of an animal is reflected through the blood parameters. So the studies on haematological parameters are important for monitoring the fish health conditions. Phosphamidon impairs metabolic activities through tissue damage leading to reduced growth and health of organism. (Tripathi G., and S. P. Shukla 1988; Sailajaraghuram D., and B.N., Naidu 1989; Khangarot, B.S., et al., 1991; Reddy D.C., et al., 1992; Mukherji M. K. and S.K., Khonr 1984 ; Mathur K.K., 1991; Sastry K.V., and A., gupta 1994; Pulla et al., Mishra 1997) Hence it become clear again that haematological investigations are greatly helpful to provide the important informations about fish health. Stein J. E., et al., (1992); De La Torre et al., (2000) reported that the blood parameters of diagnostic importance are erythrocytes and leucocytes count. Haemoglobin and haematocrit are also responded to environmental factors such as toxicological stress due to water contaminants. Numerous workers have been reported the altered haematological and biochemical parameters to make a decision for effected biological activities and the harmful consequence of contamination due to phosphamidon and

other organophosphorus pesticides (Goel et al., 1986; Khangarot B.S., and P.K., Ray 1988e; Gupta. A. K., 1995; Poonmani R., and B., Dhanakkodi 1996; Jeba kumar S.R.D. et al., 1997; Sastry et al., 1997; Imbamani N., and R., Shrivassan 1998; Rajamannar K., and L., Manohar. 1998; Lata et al., 2001; Patnail et al., 2002).

The haematological and biochemical data's are presented in table 09 and 10 separately for acute and chronic toxicity tests. The table shows that the reduction was originated in values of haemoglobin percentage (Hb%), TEC and PCV behind every part of exposure periods to lethal and sub lethal concentrations of phosphamidon. The decreased response in figures of Hb% and PCV were positively correlated with TEC. The anemic response may possible due to the inhibition of erythrocytes production. (Shekhar P., and I., Christy 1996; Anand Kumar, A. and Tripathy N.K. 2001; Chandra et al., 2001) Haemoglobin is synthesized in the body of an organism from protein and iron. Liver is the store house of iron in the body. Phosphamidon is known to bring about necrotic changes in the liver of the fishes (Verma et al., 1981). This might has led to non availability of enough iron in the body of exposed fishes leading to a decrease in Hb% which in turn decreases the oxygen carrying capacity of the blood (Chandshekhar S., and N. Jayabalan 1993; Anand Kumar, A. and Tripathy N.K. 2001). The behavioural responses also confirm the low O₂ consumption.

Table 09 and 10 shows that the levels of TLC and ESR increased after all exposure periods of phosphamidon. The level of TLC increased because leucocytes are involved in the regulations of immunological function of body (Santa kumar et al., 2000). An

increased in TLC thus occurs as a protective response to stress (Das R., 1998). Hence it become clear that TLC and ESR have negative correlation with TEC. (Goel K A., and Maya 1986; Kumar B., Benerjee V., 1990; Singh S. and Bhati D.P.S., 1991). Increased level in the values of ESR was also observed by Gupta et al., (1995); Singh and Bhati (1991); Mala F.A., G. Sharma (2009)..

The reduction in the values of Hb%, TEC and PCV indicates that the fishes suffer from anaemia or heamodilution. The alterations in MCV, MCH, and MCHC were also observed in comparison to control fishes, because they are interrelated with haemoglobin, TEC and PCV. Gradually falling in values of MCV, MCH and increased MCHC values were observed in the present study during acute and chronic toxicity conditions. These alterations verify that the fishes are suffering from anaemia during exposure periods due to iron deficiency and damaged tissue conditions especially liver. Similar haematological effects of phosphamidon in fishes, after various exposure periods have also been reported by many bio researchers (Chaudhry D.K., et al., 1984; Anand Kumar A., and Tripathy N.K., 2001; Bhatia et al., 2002; Das B. K., and S.C., Mukherjee 2003).

The biochemical parameters (blood glucose) is also an important indicator to study the internal environment of fishes. A significant increase was observed in blood glucose level intoxication of phosphamidon after acute and chronic exposure periods (Table 09 and 10). The change in blood glucose may be due to stress in fishes (Nataragan G.M., 1989; Shobha R.J.V., et al., 1989; Chandrasekhar and Jayabalan S., 1990; Sikoki, F. D., & Enajekpo, H.O.S., 1991; Chandshekhar S. and N., Jayabalan 1993; Das B.K., and S.K.,

Mukharjee 2000; Luskova et al., 2002; Saufy et al., 2007). The stress related hyperglycemia was reported in many species of teleost. It was noticed that cortisol has shown to promote catabolism of peripheral tissue via gluconeogenesis, leading to hyperglycemia. Increase in blood glucose might have resulted from gluconeogenesis to provide energy for the increased metabolic demands imposed by insecticides stress (Ghosh T. K., 1989; Jyothi B., and G., Narayan 1999; Kumar Hemant and Gupta A.B., 1997).

The depletion of glycogen content in liver and muscles by the effect of pesticides on fishes was also observed by many workers (Srivastava A.K., 1981; Nemcsok and Bones 1982; Singh H.H., Srivastava A. K. 1982; Nataragan G. M., 1989; Luskova V., et al., 2002; Medda et al., 2003). The impaired carbohydrate metabolism may also causes hyperglycemia as shown in the present study both in acute and chronic exposure of phosphamidon (Chandshekhar S. and N., Jayabalan 1993 Kumar Hemant and Gupta A.B., 1997).

(c) Study of seasonal variation:-

The data obtained in figure 23-31 and 32-40 showed that haemoglobin percentage, TEC and PCV were decreased but TLC, ESR and blood glucose level were increased following exposure to phosphamidon from untrited fishes both during acute and chronic experiment. The maximum values of Hb%, TEC, TLC, and PCV were observed at the end of rainy season and lowest values of parameters were found in strong summer season. The average values of all the parameters in control fishes were found in winter season. This could be due to the process of haemoconcentration occurs during the late

spring and summer followed by haemodilution in the late summer. The highest blood glucose level was found in rainy season. It indicates that the low temperature and rich food supply stimulate metabolism in fish body. The values of MCV, MCH and MCHC were also fluctuated according to the levels of Hb%, TEC, and PCV. The lower activity of all the parameters in summer season may also be due to the high temperature of water as shown in chapter 4, O₂ availability was also reduced in summer season (Schaefer A.A., 1925; Kaplan and Carousa 1956; Raizada et al., 1983; Harding J., and Hogland L. B., 1984; Joshi P.C., 1989; Denton J.E., and Yousef M.K., 1995; Collazos et al., 1998) causes the decreased activity of TEC, HB%, and PCV.

Determination of LC₅₀ for Phosphamidon on *Channa punctatus*

Table 06: First exploratory test

S.No.	Conc. ml/liter	No. of fishes	24 Hours		48 Hours		72 Hours		96 Hours	
			M	%M	M	%M	M	%M	M	%M
1.	0.002	5	0	0	0	0	0	0	0	0
2.	0.040	5	5	100	-	-	-	-	-	-

Table 07: Second exploratory test

S.No.	Conc. ml/liter	No. of fishes	24 Hours		48 Hours		72 Hours		96 Hours	
			M	%M	M	%M	M	%M	M	%M
1.	0.007	5	0	0	0	0	0	0	1	20
2.	0.014	5	0	0	1	20	1	40	2	80
3.	0.021	5	2	40	2	80	1	100	-	-
4.	0.028	5	4	80	1	100	-	-	-	-

Table 08: Definitive test

S.No.	Conc. ml/liter	No. of fishes	24 Hours		48 Hours		72 Hours		96 Hours	
			M	%M	M	%M	M	%M	M	%M
1.	0.008	10	0	0	0	0	1	10	2	30
2.	0.010	10	0	0	0	0	2	20	2	40
3.	0.012	10	0	0	1	10	2	30	3	60
4.	0.014	10	0	0	2	20	2	40	3	70
5.	0.016	10	1	10	2	30	3	60	3	90
6.	0.018	10	2	20	2	40	3	70	3	100
7.	0.020	10	3	30	4	60	3	90	1	100
8.	0.022	10	4	40	4	80	2	100	-	-
9.	0.024	10	6	60	4	100	-	-	-	-




Fig 21: LC₅₀ of Phosphamidon after 24 hours and 48 hours




Fig 22: LC₅₀ of Phosphamidon after 72 hours and 96 hours

Table No. 09

Effect of acute toxicity of Phosphamidon on selected blood parameters
in fresh water fish *Channa punctatus*

S No.	Parameters	Control	Exposure Period			
			24 Hours	48 Hours	72 Hours	96 Hours
01.	Hb (g/100ml)	14.1 ±0.72	10.9** ±0.63	11.5** ±0.10	11.1** ±0.49	11.5** ±0.36
02.	TEC $\times 10^6/\text{mm}^3$	3.85 ±0.07	3.42** ±0.05	3.45** ±0.07	3.48 ±0.03	3.49** ±0.04
03.	TLC $\times 10^3/\text{mm}^3$	3.2 ±0.32	4.0* ±0.20	4.2* ±0.47	4.2* ±0.40	4.3* ±0.65
04.	ESR (mm)	2.5 ±0.11	3.9 ±0.15	4.1 ±0.34	4.1 ±0.23	3.8* ±0.55
05.	PCV%	33.33 ±3.05	22.33* ±3.05	24.33* ±3.50	25.66* ±3.51	26.66* ±3.21
06.	MCH pg	36.66 ±1.34	31.55* ±1.53	33.30* ±0.43	31.26* ±1.37	33.10* ±1.03
07.	MCHC %	42.51 ±2.06	49.33* ±4.24	47.80* ±6.60	44.11* ±7.10	43.56* ±5.54
08.	MCV μm^3	86.82 ±7.86	64.45* ±10.65	70.57* ±9.73	75.85* ±0.48	76.86* ±9.63
09.	Glucose (Units)	64.73 ±5.60	70.07* ±6.53	73.71* ±6.45	73.12* ±4.67	74.91* ±4.93

* - Significant at $P < 0.01$; ** - Significant at $P < 0.001$

Table No. 10

Effect of chronic toxicity of Phosphamidon on selected blood parameters in fresh water fish *Channa punctatus*

S No.	Parameters	15 Days		30 Days		45 Days	
		C	T	C	T	C	T
01.	Hb (g/100ml)	13.9 ±0.60	11.9** ±0.36	14.6 ±0.57	12.5* ±0.05	14.3 ±0.66	12.9 ±0.15
02.	TECx10 ⁶ /mm ³	3.77 ±0.11	3.50* ±0.09	3.80 ±0.90	3.57* ±0.08	3.88 ±0.09	3.63* ±0.08
03.	TLCx10 ³ /mm ³	3.1 ±0.40	4.0* ±0.37	3.2 ±0.35	4.2 * ±0.35	3.5 ±0.40	4.8** ±0.10
04.	ESR (mm)	2.3 ±0.20	3.6 ** ±0.11	2.3 ±0.15	3.8* ±0.75	2.4 ±0.55	4.0 ±0.62
05.	PCV%	27.66 ±2.88	19.66* ±2.08	29.66 ±3.05	21.00* ±2.64	33.00 ±2.00	23.33** ±2.30
06.	MCH pg	36.82 ±0.48	34.93* ±1.68	36.97 ±0.51	35.08* ±0.94	36.92 ±0.90	35.63* ±0.88
07.	MCHC %	49.89 ±4.56	60.82* ±4.67	47.76 ±5.43	60.29* ±7.36	43.48 ±1.89	55.80* ±5.70
08.	MCV um ³	73.34 ±7.61	52.75* ±7.40	75.39 ±9.72	58.75* ±7.16	85.04 ±4.69	64.25* ±5.68
09.	Glucose (Units)	60.46 ±8.35	70.87* ±9.14	63.55 ±8.67	76.03* ±9.48	66.03 ±5.98	76.92* ±9.47

* - Significant at P < 0.01; ** - Significant at P < 0.001

Graphical comparison of seasonal variation in acute toxicity experiment




Figure 23. Haemoglobin




Figure 24. TEC




Figure 25. TLC




Figure 26. ESR




Figure 27. PCV




Figure 28. MCH




Figure 29. MCHC




Figure 30. MCV




Figure 31. Glucose

Graphical comparison of seasonal variation in chronic toxicity experiment




Figure 32. Haemoglobin




Figure 33. TEC




Figure 34. TLC





Figure 35. ESR



36. PCV




Figure 37. MCH




Figure 38. MCHC




Figure 39. MCV




Figure 40. Blood glucose

Chapter - 7

**EFFECT OF METASYSTOX
ON BLOOD PARAMETERS OF
*CHANNA PUNCTATUS***

EFFECT OF METASYSTOX ON BLOOD PARAMETERS OF *CHANNA PUNCTATUS*

Metasystox 25% E.C. was a systemic organophosphorus insecticides used against many insects. It is a water emulsifier insecticide. The chemical composition of this compound is Oxydemeton-methyl, *S*-[2-(ethylsulfinyl) ethyl] *O, O* dimethyl. It is manufactured by Bayer India limited, Mumbai.

Results

(a) Acute toxicity bioassay:-

To find out the accurate determination of LC_{50} values, two exploratory and a definitive test were conducted. In first exploratory test 100% mortality was observed in 0.050 ml/l of metasystox while no mortality in 0.010 ml/l (table 11). In second exploratory test four ascending series of concentrations (0.015, 0.025, 0.035, and 0.045 ml/liter) were taken to get narrow range of concentrations (table 12). In second range finding test 100 % mortality occurred at 0.045 ml/l after 24 hours exposure period but 20 % mortality was seen at 0.015 after 96 hours exposure of metasystox. Then seven concentrations were selected for definitive test (table 13) on the basis of second exploratory test and record the percent mortality of each selected concentration. Now a curve was plotted between percent mortality and concentrations of metasystox and sketch a line intersecting through the concentrations and 50% mortality level (fig no. 41 and 42). The LC_{50} values determined by graphical method were found to be 0.034, 0.030, 0.026 and 0.022 ml/l for metasystox at 24, 48, 72 and 96 hours.

The effect of metasystox on fish behaviour has also been observed in *Channa punctatus*. Initially the fish gives the impression of restlessness, lethargy, rolling movement and lost their balance and natural behaviour. After some time they showed erratic and burst swimming with augmentation in respiratory frequency swimming near the water surface for oxygen uptake. Toxicant exposed fish also secretes excessive mucous in comparison to control fishes. Under higher concentrations they showed hemorrhagic patches on their skin.

(b) Haematological & biochemical study:-

The haematological & biochemical parameters of control and metasystox exposed fishes at LC_{50} concentrations are presented in table 14. The Haemoglobin percentage and total erythrocytes counts (TEC) were increased significantly after 24, 48, 72 and 96 hours of exposure periods at $P < 0.01$ and $P < 0.001$. The haemoglobin percentage in control fishes was 14.1 gm% where as in treated fishes it was 16.1, 16.2, 16.3 and 16.5 gm% at the end of 24, 48, 72 and 96 hours respectively. The Hb % level was low at 24 hours of LC_{50} and high in 96 hours exposure. Although TEC was also more than control fishes but it was gradually increased with increasing exposure periods. A decreased level was observed in TLC and ESR. The difference was statistically significant at $P < 0.01$ and $P < 0.001$. The ESR was lower in 24 hours but it was also gradually increased with running exposure periods.

The values of PCV, MCH and MCV were significantly increased at $P < 0.01$ after 24, 48, 72 and 96 hours exposures. The values of PCV and MCV for 96 hours were also significant at $P < 0.001$ and same

case was found in value of MCH during 72 hours exposure period. In case of MCHC, it was decreased significantly at $P < 0.01$. The table 14 shows that the levels of glucose increased significantly at $P < 0.01$ in treated fishes. The value of blood glucose was 63.46 in control fishes where as fishes exposed to acute toxicity concentrations of metasystox showed 68.36, 73.05, 74.58 and 76.14 units at 24, 48, 72 and 96 hours respectively.

The haematological & biochemical parameters of control and metasystox exposed fishes at 1/10 of 96 hours LC_{50} concentrations are presented in table 15. The level of Hb%, TEC, PCV, MCH and MCV were significantly increased at $P < 0.01$ in treated fishes than untreated ones after the end of 15 days, 30 days and 45 days intoxication of metasystox. The level of TEC, PCV and MCV were lesser during 15 days exposure but gradually increased with increasing exposure periods. A significant decreased level of TLC, ESR and MCHC were observed at $P < 0.01$ and $P < 0.001$. The table 15 also indicates the change in blood glucose level in *Channa punctatus*. The blood glucose level was increased significantly at $P < 0.01$ in the insecticidal treated fishes.

(C) Study of seasonal variation:-

The seasonal changes of certain haematological and biochemical parameters (glucose levels) are presented in the figure 43-51 during acute toxicity bioassay and chronic toxicity bioassay was presented in fig 52-60. In toxicity test TEC and PCV were decreased during warm season in unexposed fishes. The Hb%, TEC, and MCH were lower in late summer season, while higher in beginning of rainy

season in control fishes. The value of TLC and ESR were slightly altered during summer and rainy seasons. Although Hb% TEC, PCV, MCH and MCV increased and other parameters were decreased in treated fishes as shown in table 14, but the activity levels of all the parameters were low in summer season and high in rainy season. In case of PCV maximum value was observed in winter season in control fishes. However PCV in treated fishes was almost similar during rainy and winter season but average in summer season. The levels of MCH, MCHC and MCV were also observed. They were also fluctuated during different seasons according to the values of TEC, Hb% and PCV. The highest blood glucose levels were observed during rainy season. In chronic experiments Hb%, TEC, TLC and ESR were slightly low in late summer season. In treated fishes the levels of Hb%, TEC and PCV were moderately change and low in summer seasons comparably. The levels of MCH, MCHC and MCV were also observed during chronic exposure. They were also rise and fall during different seasons according to the standards of TEC, Hb% and PCV. The blood glucose level was higher in rainy season in control fishes while in treated fishes the highest value was observed after 45 days of exposure period.

Discussion

(a) Acute toxicity:-

Indiscriminate use of insecticides has elevated the risk of contamination of environment and aquatic habitat. Various bioresearchers have been reported the LC_{50} values of metasystox in

different fish species. Natrajan G.M., (1983) notified the LC_{50} value 0.5 mg/l at 48 hours in *Channa striatus*, while Jhon et al., (1989) reported 12.9 ppm LC_{50} for 96 hours in *Heteropneustes fossilis*. Several workers have also been reported the toxicity of metasystox and other organophosphorous insecticides (Mahajan and Juneja 1979; Dhilon and Gupta 1983; Natrajan G.M., 1983; Ghosh and Chatterji 1989; Valicre E.J. and C.J., Stickney 1989; Meenaxi Das and Sambhu Prasad 1996).

Kumar Hemant and A. B., Gupta (1997) estimated the LC_{50} values in *Heteropneustes fossilis*. According to them the LC_{50} were 29.5, 24.50, 17.00 and 17.00 ppm for 24, 48, 72 and 96 hours respectively. In the present study the LC_{50} values were 0.034, 0.030, 0.026 and 0.022 ml/liter at 24, 48, 72 and 96 hours respectively, which correlates the LC_{50} values of other workers. A review of literature showed that, aquatic environment is polluted by the use of metasystox so the present study is aimed at investigating the effect of metasystox on blood parameters of affected organisms like fishes of Bundelkhand region.

Numerous abnormal behaviours such as impression of restlessness, rolling movement and lost their natural behavior and balance, erratic swimming with increment in respiratory frequency, swimming near the water surface for oxygen uptake observed after exposure period. Comparable changes in behavior of fresh water fishes on exposure to different organophosphorus toxicants were reported by several biologists. (Salim Mustfa and Ajmal Murad 1983; Kumar Hemant and A.B., Gupta 1997; Roopa Methew et al., 1997; Sadhu et al., 2001; Vatukuru S.S., 2005). These changes could be due

to the effect of metasystox on the central nervous system. Increased opercular movement of fishes when introduced in toxic media implied rise in oxygen consumption. This also may be due to hyper excitability which involves energy expenditure and producing greater demand of oxygen. Behavioral changes resulting in restlessness and loss of balance erratic swimming and hemorrhage have also been observed in the fishes by other workers (Cleveland et al., 1972; Jai Nath and Rao 1984; Beitinger T. L., 1990; Little E. E., and S. E. Finger 1990; Little et al., 1993 b; Allin C.J., and R.W., Wilson. 2000; Brewer et al., 2001; Singh Snehlata, Sadhu D. N., 2001; Vatukuru S. S., 2005).

(b) Haematological and biochemical study:-

Organophosphorus insecticides have been widely used by farmers to control insects, pests and disease vectors. They ultimately find their way into aquatic habitats such as river, lakes and ponds. They become highly toxic and ultimately disturb the food chain. The haematological and biochemical studies are useful in assessing the health of fish subjected to changing in environmental conditions. The pesticides play an important role to alter the blood parameters of fishes to a considerable extent. Keeping this in view the study was conducted to assess to the changes in haematological and biochemical parameters in fishes (*Channa punctatus*) exposed to lethal and sub lethal concentrations of metasystox. Many workers have been contributed the study of haematological changes due to organophosphorus pesticides in fishes (Agrawal S. J., and A. K., Srivastava 1980; Jagdish Mishra and Anil K., Shrivastava 1981; Singh B. B. & Narain, A. D., 1982 Pandey et al., 1980; Balint et al., 1995; Edsall, C. C., 1999; Nath. Ravindra and Banerajee V., 1999).

Haematological data are presented in table 14 and 15. Elevations in Hb%, TEC, PCV were significant at $P < 0.01$ and $P < 0.001$ after the end of all exposure period. Several bioresarches have mentioned the similar reports of Hb%, TEC and PCV using metasystox during acute and chronic toxicity exposure (Mahajan and Juneja 1979; Dhilon and Gupta 1983; Natrajan G.M., 1983; Valicre E.J., and C.J., Stickney 1989; Meenaxi Das and Sambhu Prasad 1996; Josiph P., Jhon 2007). In the addition metasystox can cause the low pH value during and chronic exposure period, so that the increased value of Hb%, TEC and PCV may be possible due to low pH value (Valicere E.J., 1989).

In the study with brook trout exposure to sub lethal conc. of cupric chloride also found significant increased in Hb% and PCV content Mitchell (1987) and Natrajan G. M., (1983) also showed the increased level of Hb%, TEC and PCV. It has been suggested that these haematological disturbances are a haemopoietic or erythrocytic mobilisation response caused by stress (Goel, K. A., 1982; Adams S.M., 1990; Barton B.A., and G.K., Iwama 1991; Wilson R.W., Taylor, E.W., 1993; Stein et al., 1992; Mikko Nikinmaa and Virpi Tervonen 2003; Vander et al., 2003).

In many species of fishes, the spleen is the primary storage site of erythrocytes (Gallaugher and Farrell 1998). Liberation of erythrocytes from the spleen is under adrenergic nervous or hormonal control. Thus, erythrocyte liberation can occur very rapidly, and is known to take place as a response to both exercise and hypoxic condition (Mahajan and Juneja 1979; Yamamoto et al. 1980, Yamamoto et al. 1983). The pioneering study by Swift and Lloyd (1974) showed that in acute hypoxia the urine flow rate of rainbow

trout increased simultaneously with an increase in blood haematocrit value. The increased excretion of water causes a reduction in plasma volume, as is observed in rainbow trout (Mahajan and Juneja 1979; Valicere E.J., 1989; Mikko Nikinmaa and Virpi Tervonen 2003). A decrease in plasma volume will increase the number of erythrocytes per unit volume.

The level of TLC were decreased after acute and chronic toxicity bioassay agrees with the report, that the release of epinephrine during stress condition causes a decrease in the number of leucocytes counts, which show the weakening process of immune system (Alkaham et al., 1998; Ajani F., 2008; Khalid et al., 2008; Natrajan G.M., 1983).

Alkahem et al., (1998) and Ajani F., (2008) also showed the decreased no. of leucocytes after being exposed to organophosphorus pesticides. This is attributed to the reduction in the no. of lymphocytes Maule and Schreer (1990) also noted the stress and level of cortisol affect the number of lymphocytes in immune organs.

In present study increased level of MCH, and MCV were observed while MCHC were decreased after the end of all exposure periods with metasystox. These alterations confirm polycythaemia in fishes due to stress condition (Goel K.A., 1982; Khattak I.U.D., Natrajan G.M., 1983). Another reason in alterations in haematological parameters in the lethal and sub lethal concentration of metasystox apper to due to damage respiratory epithelium. Mucus is also lost from the gills which causes the adverse effect in the absorption of O₂ (Natrajan G.M., 1983). Similar observation in depletion of O₂

consumption was made by Rath S., and Mishra B.N. (1980). Inhibited respiration was also observed by other workers using other organophosphorus insecticides (Hiltibran R. C. 1974; Kawatski J. A., and Mc. Donald M. J. 1974; Ranke B., and Rybicki 1975; Rath S., and Mishra B.N. 1980). The low rate of O₂ consumption caused hypoxic condition which in turn increase Hb% and TEC.

To study the health of fishes biochemical parameters are important indicators. (Jagdish Mishra and Anil K., Shrivastava 1981; Mishra, J., Shrivastava A. K., 1983; Edsall C.C., 1999; Vdesek J., et al., 2009). Changes in the biochemical parameters cause changes in metabolism due to the effect of various toxicants. Changes in the activity of several enzymes and carbohydrate metabolism have been observed by several workers (Hochachka et al., 1978; Shrivastava A.K., 1981; Nemcsok and Bones 1982; Singh H.H., Shrivastava A. K. 1982; Ghosh T.K. ,1989; Natarajan G. M., 1989; Kumar Hemant and Gupta A.B.1997; Luskova V., et al., 2002; Saufy H., et al., 2007). The blood glucose level in Indian cat fish *Heteropneustes fossilis* exposed to sub lethal concentrations of other toxicant was also found to increase considerably (Gill T.S., et al., 1990; Ceron, J.J et al., 1997; Kumar Hemant and Gupta A.B. 1997; Shrivastva and Singh, 1981; Saufy H., et al., 2007). The blood glucose level was increased significantly at $P < 0.01$ intoxication of metasystox after end of acute and chronic exposure periods during this toxicological investigation. It may be due to disturbing process of impulse across the neuromuscular neural junctions by inhibiting the enzymes AChE, which modulate the amount of neurotransmitter acetylcholine. The hyperglycemic conditions induced by pesticides might be explained in the part of

inhibition of cholinesterase at neuroeffectors sites in the adrenal medulla leading to hypersecretions of adrenaline, which stimulate the break down of glycogen into glucose (Hochachka et al., 1978; Shastri K.V., and Siddiqui A. A., 1982; Ghosh T.K., 1987; Ghosh, T. K., 1989; Jyothi B., and Natrajan G. M., 1999; Kumar Hemant and Gupta A.B., 1997).

(c) Study of seasonal variation:-

The data presented in figure 43-51 showed that in acute toxicity test Hb%, TLC, ESR and MCH were lower in late summer season, while higher in beginning of rainy season in control fishes. The value of Hb% was slightly altered during summer and rainy seasons. This is because of haemodilution at the onset of rainy season. The lower activity of all parameters in the summer season may also be due to the high temperature of water. O₂ availability was also reduced in summer season. Although all the activity level were similar as shown in table 14 and 15 such as some parameters were increased and some were decreased with corresponding control fishes but the level was low during summer season and high in rainy season. It was also noticed that fish differ in their tolerance limit to temperature, dissolved oxygen (DO), pH, alkalinity, and other pollution factors. Mainly the temperature fluctuations affect the health of fishes. When water is extremely heated, much energy, oxygen and vapour is released into the air, leaving behind a high concentration of CO₂, which makes the water more acidic which increases the haemodilution. Wademeyer and Yasutake (1977) listed a number of environmental and physiological variables which show relationship to fish health. The highest blood glucose level found in rainy season in control fishes as well as the

level of glucose increase in rainy season of treated fishes indicates stimulated metabolism at low temperature and rich food supply. Several workers have also reported the various alterations in blood parameters due to different environmental stress (Charistoforides C. & Hedley-Whyte J., 1969; Bridges et al., 1976; Cech J.J., and D.E., Wohlschlag 1981; Barton B. A., and G. K. Iwama., 1991; Stein et al., 1992; Wilson R.W., Taylor E.W., 1993; Collazos et al., 1998).

Determination of LC₅₀ for Metasystox on *Channa punctatus*

Table 11: First exploratory test

S.No.	Conc.ml /liter	No. of fishes	24 Hours		48 Hours		72 Hours		96 Hours	
			M	%M	M	%M	M	%M	M	%M
1	0.01	5	0	0	0	0	0	0	0	0
2	0.05	5	5	100	-	-	-	-	-	-

Table 12 : Second exploratory test

S.No.	Conc. ml/ liter	No. of fishes	24 Hours		48 Hours		72 Hours		96 Hours	
			M	%M	M	%M	M	%M	M	%M
1	0.015	5	0	0	0	0	0	0	1	20
2	0.025	5	0	0	1	20	1	40	1	60
3	0.035	5	3	60	1	80	1	100	-	-
4	0.045	5	5	100	-	-	-	-	-	-

Table 13: Definitive test

S.No.	Conc.ml /liter	No. of fishes	24 Hours		48 Hours		72 Hours		96 Hours	
			M	%M	M	%M	M	%M	M	%M
1	0.016	10	0	0	0	0	1	10	2	30
2	0.02	10	0	0	1	10	1	20	2	40
3	0.024	10	1	10	1	20	2	40	2	60
4	0.028	10	2	20	2	40	2	60	3	90
5	0.032	10	3	30	3	60	3	90	1	100
6	0.036	10	7	70	2	90	1	100	-	-
7	0.04	10	9	90	1	100	-	-	-	-



Fig 41: LC₅₀ of Metasystox after 24 hours and 48Fig 42: LC₅₀ of Metasystox after 72 hours and 96 hours

Table No. 14

Effect of acute toxicity of Metasystox on selected blood parameters in
fresh water fish *Channa punctatus*

S No.	Parameters	Control	Exposure Period			
			24 Hours	48 Hours	72 Hours	96 Hours
01.	Hb (g/100ml)	14.10 ±0.74	16.1* ±0.81	16.2* ±0.77	16.3* ±0.65	16.5** ±0.35
02.	TEC $\times 10^6/\text{mm}^3$	3.77 ±0.10	3.87* ±0.02	3.89* ±0.05	3.91* ±0.04	3.94* ±0.02
03.	TLC $\times 10^3/\text{mm}^3$	3.3 ±0.17	2.3** ±0.11	2.4** ±0.15	2.5** ±0.15	2.7* ±0.20
04.	ESR (mm)	2.1 ±0.36	0.6** ±0.28	0.7* ±0.46	0.8* ±0.51	1.0** ±0.11
05.	PCV%	33.33 ±3.21	40.33* ±2.88	42.00* ±2.64	43.66* ±3.21	47.00** ±1.73
06.	MCH pg	37.08 ±1.01	41.50* ±2.35	41.77* ±1.98	41.64** ±1.24	39.84* ±3.54
07.	MCHC %	44.69 ±4.62	39.96* ±1.14	39.35* ±0.11	37.04* ±1.68	35.19* ±0.81
08.	MCV μm^3	83.72 ±11.04	103.97* ±7.90	107.86* ±6.69	111.54* ±7.29	119.07** ±3.98
09.	Glucose (Units)	63.46 ±4.83	68.36* ±5.11	73.05* ±5.88	74.58* ±4.70	76.14* ±5.07

* - Significant at $P < 0.01$; ** - Significant at $P < 0.001$

Table No. 15

Effect of chronic toxicity of Metasystox on selected blood parameters
in fresh water fish *Channa punctatus*

S No.	Parameters	15 Days		30 Days		45 Days	
		C	T	C	T	C	T
01.	Hb (g/100ml)	13.6 ±0.58	15.8* ±0.60	13.8 ±0.55	15.6* ±0.72	14.0 ±0.56	15.4* ±0.72
02.	TECx10 ⁶ /mm ³	3.71 ±0.11	4.01* ±0.14	3.74 ±0.12	4.03* ±0.13	3.76 ±0.13	4.06* ±0.13
03.	TLCx10 ³ /mm ³	3.3 ±0.40	2.4 ** ±0.37	3.6 ±0.35	2.6** ±0.35	3.4 ±0.40	2.7** ±0.10
04.	ESR (mm)	2.4 ±0.36	1.1** ±0.17	1.8 ±0.23	0.9* ±0.36	2.2 ±0.26	0.9* ±0.40
05.	PCV%	27.33 ±2.30	34.33* ±2.08	28.66 ±2.88	36.66* ±2.30	30.00 ±2.64	37.33* ±2.88
06.	MCH pg	36.82 ±0.61	39.57* ±0.93	35.86 ±2.05	39.16 * ±0.44	37.28 ±0.79	39.61* ±2.39
07.	MCHC %	46.92 ±4.00	41.14* ±2.75	48.42 ±4.78	36.57* ±1.19	40.57 ±4.84	40.56* ±2.39
08.	MCV um ³	73.91 ±5.52	89.58* ±8.45	76.52 ±6.18	92.86* ±4.46	83.30 ±2.26	97.78 ** ±1.47
09.	Glucose (Units)	61.55 ±5.01	69.93* ±5.59	64.86 ±4.99	71.35* ±5.83	65.61 ±4.11	74.85* ±4.45

* - Significant at P < 0.01; ** - Significant at P < 0.001

Graphical comparison of seasonal variation in acute toxicity experiment




Figure 43. Haemoglobin




Figure 44. TEC




Figure 45. TLC




Figure 46. ESR




Figure 47. PCV




Figure 48. MCH




Figure 49. MCHC




Figure 50. MCV




Figure 51. Blood glucose

Graphical comparison of seasonal variation in chronic toxicity experiment




Figure 52. Haemoglobin




Figure 53. TEC




Figure 54. TLC




Figure 55. ESR




Figure 56. PCV




Figure 57. MCH




Figure 58. MCHC




Figure 59. MCV




Figure 60. Blood glucose

Chapter - 8

**EFFECT OF IMIDACLOPRID
ON BLOOD PARAMETERS OF
*CHANNA PUNCTATUS***

EFFECT OF IMIDACLOPRID ON BLOOD PARAMETERS OF *CHANNA PUNCTATUS*

Imidacloprid 17.8% S.L. (trade name Anumida) insecticide was used in this investigation. The chemical composition of this compound is (1- [6- chloro -3- pyridylmethyl]- N- nitroimidazolidin -2-ylideaneamine). It is manufactured by Anu products limited, Trigon road, Faridabad. India.

Results

(a) Acute toxicity bioassay:-

The LC_{50} of imidacloprid was calculated by same procedure as described earlier. In the first exploratory test 100% mortality was observed in 0.075 ml/l of imidacloprid while no mortality occurred in 0.025 ml/l. (Table 16). In second exploratory test mortality due to different concentrations were observed (Table 17). The concentrations were 0.030, 0.040, 0.050, 0.060 and 0.070 ml/liter. The 100% mortality was observed at 0.050 ml/l at 96 hours, 0.060 ml/l at 72 hours and 0.070 ml/l at 24 hours. In definitive test nine ascending series of concentrations viz 0.032, 0.036, 0.040, 0.044, 0.048, 0.052, 0.056, 0.060 and 0.064 ml/l were taken and mortality data was recorded in table 18. Then a curve was plotted between percent mortality and concentrations of imidacloprid which were used in definitive test and LC_{50} values were determined by drawing a line, intersecting through the concentration of imidacloprid at 50% mortality level (Fig 61 and 62). The LC_{50} values came to be as:-

(1.) 24 hours-0.058 ml/liter

gradually increased with increasing exposure periods. In case of ESR the value of 48 and 96 hours exposure period were also significant at $P < 0.001$. It also shows increasing trend. The levels of MCH and MCV were decreased significantly at $P < 0.01$ when compared to control fishes. While the MCHC was increased at the end of 24, 48 & 96 hours exposure periods except at 72 hours. Regarding the biochemical alterations, the glucose level increased significantly at $P < 0.01$. The level of glucose was 63.36 in untreated fishes where as fishes exposed to acute toxicity concentrations of imidacloprid showed 70.43, 71.26, 72.01, 75.19, mg/100 ml blood of glucose concentration.

In chronic toxicity bioassay (table 20) the Hb%, TEC, PCV and MCH decreased significantly ($P < 0.01$) at 15 days and 45 days after exposure of imidacloprid (1/10 of 96 hours LC_{50} concentration). The table 20 shows that the level of TEC was decreased but no significant difference were observed after the end of 15, 30 and 45 days of exposure periods. A statistically insignificant increased was observed in erythrocytes sedimentation rate expect 15 days exposure period which was significant at $P < 0.01$. TLC increased significantly during toxication. The level of MCV in treated fishes than untreated was also decreased. The level of blood glucose was increased significantly ($P < 0.01$) in treated fishes. The level of blood glucose was lower in 15 days exposure but gradually increased after the end of 30 days and 45 days exposure periods.

(c) Study of seasonal variation:-

Seasonal changes of some haematological and biochemical parameters (glucose levels) are presented in the figure 63-71 during

(2.) 48 hours-0.050 ml/liter

(3.) 72 hours-0.042 ml/liter

(4.) 96 hours-0.034 ml/liter

When the fishes were introduced into test medium with different concentrations of imidacloprid, they showed changes in their behaviour. The copious secretion of mucus by the body surface of the fishes was seen. The loss of scales, decoloration, rolling movement and loss of equilibrium in higher concentrations were observed in fishes. The fishes exhibited lethargy and erratic swimming with increment in respiratory rate. At the time of death transient hyperactivity was also observed.

(b) Haematological & biochemical study:-

The haematological and bio chemical parameters of control and Imidacloprid exposed fishes at LC_{50} concentrations are tabulated (table no. 19). The haemoglobin percentage was decreased at 24, 48, 72 and 96 hours. The difference was statistically significant at $P < 0.001$ but at 72 hours it was significant at $P < 0.01$. The haemoglobin percentage in control fishes it was 13.9 gm % where in treated fishes was 11.1, 11.4, 11.5 and 11.9 gm % for 24, 48, 72 and 96 hours of exposure periods respectively. The total erythrocytes count was also less in comparison to corresponding control. However, it was increased with increased exposure periods. Packed cell volume in control fishes was 21.66, 23.00, 23.66 and 25.33 at 24, 48, 72 and 96 hours respectively. The difference is statistically significant at $P < 0.01$. The TLC was increased significantly at $P < 0.01$. However all the values of Hb%, TEC and PCV were lower than control fishes but

acute toxicity bioassay and in figure 72-80 in chronic toxicity bioassay. The Hb%, TEC and PCV were slightly decreased during late summer season in untreated fishes. If the observation was made only in control fishes all the hematological parameters (Hb%, RBC, TEC, PCV, ESR, MCV, MCH & MCHC) and biochemical parameter (blood glucose) were decreased in summer season but slight difference was observed in winter and rainy season (fig 63-80). The haematological changes that occurred in treated fishes are similar as shown in table 19 and 20. The levels of Hb%, TEC, PCV, MCH and MCV were decreased but TLC, ESR and MCHC were increased in treated fishes when compared to untreated control during different season both in short term and long term duration as shown in fig 63-80.

Discussion

(a) Acute toxicity bioassay

The most effective insecticide against sucking insect pests currently used is imidacloprid (1- [6-chloro- 3 – pyridylmethyl] – *N* - nitroimidazolidin -2 – yldeaneamine), a neurotoxic chemical with the same mode of action as nicotine (Armbrust K.L., 2000; Matsuda et al., 2001; Tomlin 2000). It belongs to a group of insecticides referred to as the chloronicotinyl group (Kidd and James 1991; Cox et al., 1997; Tomlin 1997). It produces toxicity by binding to and over stimulating nicotinic acetylcholine (Ach) receptors on the postsynaptic membranes of neurons (Kidd and James 1991; Song et al., 1997; Tomlin 1997). Kidd and James (1991) reported LC₅₀ value 211 mg/l in rainbow trout (*Oncorhynchus mykiss*) for 96 hours. George Young (2009) reported LC₅₀ value of 237 mg/l in *Golden orfe*. The LC₅₀

values in invertebrates have also been reported by Sarah. J., (2008) which was 65.43 µg/l in *Chironomus tentans* for 96 hours. Imidacloprid has moderate toxicity to mammals. Tomlin (1997) reported the LD₅₀ in rat (*Rattus norvegicus*) which was 460 mg/kg. Compared to the available literature on imidacloprid toxicity to fishes and other aquatic invertebrates (Liu M-Y., Casida J.E., 1993; Liu et al., 1993) our results indicate that fishes are also sensitive to imidacloprid and LC₅₀ values were comparable to other workers on other animals. Very few literature are available on toxicity of imidacloprid. This study and this chemical is used as insecticide in Bundelkhand region so it become necessary to investigate the toxicity of imidacloprid. The study of imidacloprid was used for the first time in Bundelkhand region for toxicity investigation in fishes.

The behavioral changes of hyperactivity and copious secretion of mucus by the body surface of the fishes and other effects like as loss of scales, decoloration, rolling movement and loss of equilibrium in higher concentrations were observed in this study. This may be due to intoxication of imidacloprid on the central nervous system. Increased opercular movement of fishes as soon as introduced in toxic media involves considerable energy spending and there by create a greater require of oxygen. Similar changes in behavior of fresh water fishes due to exposure of other pesticides were reported by other biologists using other pesticides (Mukhopadhyay, P.K and Dehadrai, P.V. 1984; Olla et al., 1980; Salim Mustfa and Ajmal Murad 1984; Lal et al., 1984; Naqvi and Howkins 1988; Gray, R.H. 1990; Little et al., 1993b; Baldwin et al., 1994; Doving K. V., 1995; Kumar Hemant and A.B. Gupta 1997; Beauvais et al., 2000 ; Brewer, S.K., et al., 2001;

Sadhu et al., 2001; Vatukuru S.S., 2005).

(b) Haematological and biochemical study:-

In 1992, Imidacloprid was introduced as a new insecticides. It is most active against sucking insects so it is widely used in agricultural programmes. In India, renewal of agricultural programme has increased utilization of pesticides some of which are known to be highly toxic to non-target organisms such as fishes (Shrivastva, A. K. and N. N. Shrivastva. 1987; Benerjee G. and Rajendranath 1990; Benerji, G. and Rajendranath 1988; Mohapatra, B.C. and Nobal. 1992; Sastry, A. and R. A. Agrawal. 1993; Tiwari, C., 1995; Watson et al., 1997; Kumar et al., 1999; Das, R., 2000; Alam, M. N. and D. N. Sadhu, 2001). However, this positive trend is helpful in increasing food production and improving crop health the use of pesticides has negative ramifications for fish and other aquatic life inhabiting ponds, lakes, rivers, streams. On reaching aquatic ecosystems, such chemicals can cause serious harm to fishes and other aquatic organism. Studies have shown that when the water quality is affected by toxicants, physiological changes will be reflected in the values of one or more of the haematological parameters (Van Vuren, 1986; Shrivastva, A. K. and N. N. Shrivastva. 1987; Benerji, G., and Rajendranath 1988; Barto B. A., and G. K., Iwama. 1991; Fernando M.D., and E.A., Moliner 1991; Mohapatra B.C., and Nobal. 1992;).

For this approach many Bioresearches used haematological and biochemical parameters as biomarkers. In fishes, exposure to Imidacloprid pollutants can induce either increases or decreases in haematological levels. Many workers have been selected

haematological and biochemical parameters to determine the effected biological activities on test organism because they accept it, that the blood parameters are the best biomarkers to investigate the hazardous effect of pollution caused by various toxicants (Verma et al., 1981; Mishra et al., 1989; Sarkar 1990a; Sarkar 1990b; Sreenivasulu et al., 1991; Sadhu D.N., 1993; Wilson R.W.; Taylor E.W., 1993; Sastry A. and R. A. Agrawal. 1993; Jain R. and K.D., Mishra. 1995; Tiwari C., 1995; Kumar et al., 1997; Watson W.J., et al., 1997; Kumar S., et al., 1999; Das R., 2000; Delor et al., 2000; Alam M. N. and D. N., Sadhu 2001; Mishra et al., 2001; Singh S. and D. N., Sadhu 2001).

The Hb%, RBC, PCV%, the oxygen carrying capacity of blood significantly decreased after both acute and chronic exposure periods to imidacloprid compared to corresponding control. This may be due to the impairment oxygen supply to various tissues, thus resulting a slow metabolic rate and low energy production (Ahmad F., et al., 1995). Similar response were observed in the report of Blaxhall (1972); Wedemeyer et al., (1976); Gill and Pant (1985); Chakraporti P., (1986); Hoemechaudhuri (1986); Mishra B. K., (1993); Ravindra Nath and Benerjee V., (1999). The significant decreased in total RBC count, PCV, and Hb % also might be due to hypo stimulation to erythropoietic tissue (Lone et al., 1976). Decrease in TEC, Hb% and PCV% were also presented in the work of Mahajan and Juneja 1979; Mishra B. K., 1993 in *Channa punctatus* and *Heterpneustes fossilis*.

The alterations in haematological parameters were brought about by Imidacloprid. The decreased in Hb% was seen in the present study causing anaemia which is due to decrease synthesis of red blood cells (Morgen et al., 2000; and Nuri et al., 2003). The response of

MCH and MCV were decreased while the value of MCHC was fluctuated in acute exposure but in chronic exposure it was increased slightly. Similar response were also observed by (Janardhana Reddy et al., 1998; Verma G.P., and Pranamita Panighari 1998; Vatukaru S.S., 2005).

Increase in the TLC level also exhibited stress condition. The increase TLC can be correlated with an increase in antibodies production which helps in survival and recovery of the fishes exposed to imidacloprid (Jhosi P., Deep H., 2002; Ramesh M., and Saravanan M., 2008). An increased in TLC and ESR have negative correlation with TEC (Kumar B., and Benerjee V., 1990; Goel K A., and Maya 1986). Malla F.A.G., Sharma (2009) and Singh S., and Bhati DPS., (1991) also reported increased values of ESR were found in *Channa punctatus* intoxication of Chlorpyrifos (Chaturvedi L. D. and Agrawal K., 1993; Geol K.A., and Maya 1986; Kumar B., and Banerji V. 1990; Singh S., and Bhati DPS., 1991).

The blood glucose level was increased in treated fishes exposed to imidacloprid after both acute and chronic exposure. It was noticed that hyperglycemic condition arise in fresh water fishes following exposure to pesticides (Shrivastava and Singh 1987; Kumar Hemant and A.B., Gupta 1997). The increase in circulating glucose level is also correlated with increased concentration of the circulating acetylcholine under stressed conditions. In this study increased blood glucose might have resulted from gluconeogenesis to provide energy for the increased metabolic demands imposed by imidacloprid stress. Many other bio researchers have also been reported the hypoglycemic condition of fish intoxication of various organophosphorus pesticides

(Hochachka, P.W., 1978; Shaikh, Y.A., and P.K., Hiradher 1985; Ghosh, T. K., 1989; Shobha et al., 1989; Chandrasekhar and Jayabalan S. 1993; Gill T.S., et al., 1990 Sikoki, F. D., & Enajekpo, H.O.S. 1991; Ghosh R, Shrotri R .V., 1992; Ceron J.J., et al., 1997; Kumar Hemant and Gupta A.B., 1997).

(c) Study of Seasonal variation:-

The data obtained in figure 63-71 showed that Haemoglobin percentage, TEC, PCV, MCH and MCV decreased but TLC, ESR, MCHC and blood glucose level were increased following exposure to imidacloprid from control fishes both in acute and chronic experiment. These changes were presented in table 19 and 20, but the concentrations of above parameters in control and treated fishes were lowest in summer and then increased at the end of late summer. The increased PCV in rainy season was observed in control fishes. The changes in haematological and biological parameters may be due to low value of dissolve oxygen in summer. Hence low oxygen availability from water increases the process of haemodilution in summer followed by haemoconcentration in winter. The another reason of low blood parametric value in late summer was increasing temperature , increasing CO₂, and decreasing the oxygen affinity in the blood (Eddy F.B. 1973). The high level of CO₂ causes low ph value (chapter 4) in summer is the indication of haemodilution. The temperature of water had a direct influence on the toxicity of many pollutants on the growth of fishes. Therefore the blood parameters were low in summer season. (Akira Kakuno and Jiro Koyama 1994; Gross et al., 1996; Best et al, 2001).

Determination of LC₅₀ for Imidacloprid on *Channa punctatus*

Table 16: First exploratory test

S.No.	Conc.ml /liter	No. of fishes	24 Hours		48 Hours		72 Hours		96 Hours	
			M	%M	M	%M	M	%M	M	%M
1	0.025	5	0	0	0	0	0	0	0	0
2	0.075	5	5	100	-	-	-	-	-	-

Table 17: Second exploratory test

S.No.	Conc.m l/liter	No. of fishes	24 Hours		48 Hours		72 Hours		96 Hours	
			M	%M	M	%M	M	%M	M	%M
1	0.03	5	0	0	0	0	0	0	1	20
2	0.04	5	0	0	1	20	1	40	2	80
3	0.05	5	1	20	1	40	2	80	1	100
4	0.06	5	3	60	1	80	1	100	-	-
5	0.07	5	5	100	-	-	-	-	-	-

Table 18 : Definitive test

S.No.	Conc.m l/liter	No. of fishes	24 Hours		48 Hours		72 Hours		96 Hours	
			M	%M	M	%M	M	%M	M	%M
1	0.032	10	0	0	1	10	1	20	2	40
2	0.036	10	0	0	1	10	2	30	3	60
3	0.04	10	0	0	2	20	2	40	4	80
4	0.044	10	1	10	2	30	3	60	3	90
5	0.048	10	2	20	2	40	3	70	3	100
6	0.052	10	3	30	3	60	3	90	1	100
7	0.056	10	4	40	3	70	3	100	-	-
8	0.06	10	6	60	2	80	2	100	-	-
9	0.064	10	8	80	2	100	-	-	-	-



Fig 61: LC₅₀ of Imidacloprid after 24 hours and 48 hoursFig 62: LC₅₀ of Imidacloprid after 72 hours and 96 hours

Table No. 19

Effect of acute toxicity of Imidacloprid on selected blood parameters
in fresh water fish *Channa punctatus*

S No.	Parameters	Control	Exposure Period			
			24 Hours	48 Hours	72 Hours	96 Hours
01.	Hb (g/100ml)	13.9 ±0.60	11.1** ±0.75	11.4** ±0.35	11.5* ±1.00	11.9** ±0.20
02.	TEC $\times 10^6/\text{mm}^3$	3.77 ±0.11	3.46* ±0.05	3.48* ±0.07	3.50* ±0.06	3.52* ±0.50
03.	TLC $\times 10^3/\text{mm}^3$	3.1 ±0.40	3.6* ±0.58	4.0* ±0.41	4.1* ±0.43	4.2* ±0.70
04.	ESR (mm)	1.9 ±0.11	3.8* ±0.28	3.6** ±0.57	3.8* ±0.28	4.0** ±0.64
05.	PCV%	30.00 ±3.60	21.66* ±1.52	23.00* ±3.48	23.66* ±2.08	25.33* ±2.51
06.	MCH pg	36.85 ±0.47	32.26* ±2.63	33.09* ±1.66	33.67* ±2.11	34.10* ±1.59
07.	MCHC %	46.63 ±2.42	52.92* ±2.35	51.94* ±6.59	46.05* ±2.35	42.83* ±0.42
08.	MCV μm^3	79.43 ±7.26	62.54* ±3.99	67.96* ±5.94	67.56* ±6.29	72.11* ±6.17
09.	Glucose (Units)	63.36 ±6.29	70.43* ±5.24	71.26* ±3.48	72.01* ±6.75	75.19* ±7.05

* - Significant at $P < 0.01$; ** - Significant at $P < 0.001$

Table No. 20

Effect of chronic toxicity of Imidacloprid on selected blood parameters in fresh water fish *Channa punctatus*

S No.	Parameters	15 Days		30 Days		45 Days	
		C	T	C	T	C	T
01.	Hb (g/100ml)	13.9 ±0.60	11.5** ±0.30	14.1 ±0.55	11.8** ±0.65	14.2 ±0.56	12.1** ±0.34
02.	TEC $\times 10^6/\text{mm}^3$	3.77 ±0.11	3.17 ±0.20	3.82 ±0.09	3.24 ±0.05	3.78 ±0.25	3.31* ±0.05
03.	TLC $\times 10^3/\text{mm}^3$	3.1 ±0.40	4.0* ±0.23	3.3 ±0.30	4.3* ±0.34	3.4 ±0.20	4.3* ±0.79
04.	ESR (mm)	1.9 ±0.11	3.3** ±0.28	2.0 ±0.05	3.9 ±0.36	2.1 ±0.28	4.1 ±0.23
05.	PCV%	27.66 ±2.88	20.00* ±2.48	29.66 ±3.05	21.00* ±3.48	32.33 ±2.30	23.00* ±3.46
06.	MCH pg	36.85 ±0.47	36.34* ±1.12	36.89 ±0.52	36.58* ±1.95	37.69 ±1.10	36.47* ±0.98
07.	MCHC %	50.59 ±5.47	55.71* ±6.53	46.55 ±3.52	57.19* ±6.49	45.47 ±3.75	53.27* ±6.84
08.	MCV μm^3	73.41 ±7.73	63.07* ±11.38	77.70 ±8.24	64.73* ±10.55	85.71 ±6.82	69.33* ±10.37
09.	Glucose (Units)	59.40 ±7.58	67.82* ±8.41	63.29 ±7.84	74.17* ±8.74	64.98 ±4.67	76.23* ±7.05

* - Significant at $P < 0.01$; ** - Significant at $P < 0.001$

Graphical comparison of seasonal variation in acute toxicity experiment




Figure 63. Haemoglobin




Figure 64. TEC




Figure 65. TLC




Figure 66. ESR




Figure 67. PCV




Figure 68. MCH




Figure 69. MCHC




Figure 70. MCV




Figure 71. Blood Glucose

Graphical comparison of seasonal variation in chronic toxicity experiment




Figure 72. Haemoglobin




Figure 73. TEC




Figure 74. TLC




Figure 75. ESR




Figure 76. PCV




Figure 77. MCH




Figure 78. MCHC




Figure 79. MCV





Figure 80. Blood glucose

- 
- **SUMMARY**
 - **BIBLIOGRAPHY**
 - **ABBREVIATION AND
SYMBOLS**
 - **ANNEXURE**

SUMMARY

Aquatic toxicology is concerned with the adverse effects of various toxicants. The pesticides are mostly used for the control of insects and weeds to improve crop yield in agriculture in the area where this study was carried out. The pesticides are finally find their way into the aquatic environment through water run off. These pesticides have been reported to negative consequence on biota and the environment at large scale. The pesticides have deleterious effects not only on target organisms but also on non target organisms like fishes. The healthy fishes are much affected by these pesticides because they are highly sensitive to the pesticides. Bundelkhand has rich sources of water reservoir as Betwa river, Cane river, Matatila dam and Pahunj river. These reservoirs have become rich sources of fish farming and valuable source of fish food reaching to different areas of our country. The *Clarias batrachus*, *Labeo rohita*, *Heteropneustes fossilis* and *Channa punctatus* species are some fishes mainly used for food. These fishes are continuously challenged by toxicants used in this area. Keeping this in view four toxicants one Glyphosate (herbicide) and three oraganophosphorus pesticides viz Phosphamidon, Metasystox and Imidacloprid were selected to evaluate data on their toxicity in fresh water fishes *Channa punctatus* of Bundelkhand region. Studies have been shown that when the water quality is affected by toxicants, any physiological changes will be reflected by the haematological and biochemical parameters in fishes. So it is necessary to investigate that how much concentration is required to cause adverse effects of fishes, what blood components are affected by the chemical pesticides and what blood functions are

impaired. Therefore, the aim of this work is the determination of LC_{50} of different toxicants (Glyphosate, Phosphamidon, Metasystox and Imidacloprid) and their effects on blood parameters of *Channa punctatus*. The teleost fish *Channa punctatus* was selected in this study because of its wide availability and edibility in India. The literature survey also confers that very little work has been conducted on these toxicants. The toxicological studies in fresh water fishes *Channa punctatus* following exposure of glyphosate and imidacloprid are conducted for the first time in the present thesis.

Chlorine free tap water was used through out the course of the experiment. The physiological characters of water sample like the temperature of the test medium, dissolve oxygen, alkinity, hardness and specific conductivity were tested in the Zonal laboratory U.P. Jal Nigam Babina, Jhansi. The physiological characters of water during all three seasons rainy, summer and winter respectively were pH 7.4, 6.8 and 7.2, dissolve oxygen 7.6, 6.2 and 8.9 mg/liter, alkinity 326, 320 and 308 mg/liter as $CaCO_3$, hardness 120, 11.7 and 128.7 mg/liter as Ca, specific conductivity 792, 765 and 782 micro mho.

The fishes (*Channa punctatus*) were collected from different water bodies of Bundelkhand region with the help of professional fisher man. Live and healthy fishes were used in all the toxicological investigations. The selected fishes were checked against injury or infection by keeping 0.2% of potassium permanganate solution for 1-2 minute. They were kept in glass aquaria having a capacity of more than 40 liters. All over the experiment water was changed daily. The fishes were acclimatized in laboratory condition for 6-10 days. During acclimatization the fishes were fed egg albumin, earth worms and

small insects.

Under acute toxicity study LC_{50} values after 24h, 48h, 72h and 96h were determined by direct interpolation method, which includes two exploratory and a definitive test. The mortality was recorded after a period of 24, 48, 72 and 96 hrs and dead fishes were removed when observed. In first exploratory test (two concentrations lower and higher) were employed in jar containing five fishes each. Then four and five concentrations were taken to find out narrow range of concentrations in second exploratory test. On the basis of second range finding test 7-9 concentrations were selected for definitive test. The concentrations from the definitive test were employed to determine the LC_{50} values by plotting a dose response curve between percent mortality and concentrations of toxicants. A line was drawn between the point represent the % mortality and concentrations. The concentrations at which this line crosses for the 50% lethality line was the actual lethal concentration of toxicant. Therefore the LC_{50} values of all the selected toxicants (Glyphosate, Phosphamidon, Metasystox and Imidacloprid) were determined:-

- The LC_{50} values of Glyphosate in *Channa punctatus* conducted to be as 0.018, 0.015, 0.012 and 0.009 ml/liter at 24, 48, 72 and 96 hours.
- The LC_{50} values of Phosphamidon in *Channa punctatus* were 0.023, 0.019, 0.015 and 0.011 ml/liter at 24, 48, 72 and 96 hours.
- The LC_{50} values of Metasystox in *Channa punctatus* were found to be as 0.034, 0.030, 0.026 and 0.022 ml/liter at 24, 48,

72 and 96 hours.

- The LC_{50} values of Imidacloprid were 0.058, 0.050, 0.042 and 0.034 ml/liter at 24, 48, 72 and 96 hours in *Channa punctatus*.

After calculating the LC_{50} values 40 fishes were collected, acclimatized and divided into 5 groups (A, B, C, D and E) for acute toxicity bioassay. LC_{50} concentrations of selected toxicants were added to the groups A, B, C and D respectively. The fifth group was running without toxicant served as control. The blood was collected after 24, 48, 72 and 96 hours respectively from each group, serum was separated and haematological and biochemical parameters were tested.

For chronic toxicity test 60 fishes were collected from market and washed with 0.2% $KMnO_4$ to avoid any dermal infection and acclimatized for at least 10 days in laboratory condition. The fishes were divided into two groups of 30 each. Group one was exposed to sub lethal concentration (1/10 of 96 hours LC_{50}) of toxicants. The second group was kept as untreated control. After 15, 30 and 45 days 10 fishes from each group were sacrificed, blood was collected and serum was separated for the testing of blood parameters. The whole technique of acute and chronic toxicity bioassay were applied for all the toxicants. All the acute and chronic (blood parameters examinations) studies were also conducted in different seasons viz winter, summer and rainy season.

Glyphosate:-

In acute toxicity test Hb % and TEC were decreased significantly ($P < 0.001$). A significant decrease was also observed in

levels of PCV, MCH & MCV at ($P < 0.01$). The other parameters like TLC and MCHC levels increased significantly at ($P < 0.01$). ESR was also increased from the control fishes. The level of glucose was increased significantly ($P < 0.01$) in treated fishes.

In chronic toxicity bioassay the Hb %, TEC, PCV, MCH & MCV decreased significantly ($P < 0.01$) after 15 days, 30 days and 45 days exposure of Glyphosate. Significant increased in the level of TLC and MCHC was observed ($P < 0.01$). Although an increased level of ESR was observed when compared with the unexposed fishes but statistically the difference was not significant. Glucose level were elevated significantly ($P < 0.01$) when the fishes were subjected to glyphosate treatment.

Phosphamidon:-

The Hb % TEC were decreased significantly $P < 0.001$. Although the values of Hb% and TEC were all the times lesser than untreated fishes but at 72 hours the total erythrocytes counts was insignificant. The values of ESR were insignificantly increased, but at 96 hours it was significant ($P < 0.01$). The data illustrated that the value of PCV, MCH and MCV were significantly decreased ($P < 0.01$). TLC and MCHC were found to increase at $P < 0.01$. The glucose level estimated in treated fishes were markedly increased after all exposure periods.

In chronic toxicity bioassay the levels of Hb % TEC, PCV, MCH & MCV were decreased after 15 days, 30 days and 45 days exposure at 1/10 of 96 hours of phosphamidon LC_{50} concentration. It was observed that the levels of Hb%, TEC, MCH, PCV and MCV

were less than control fishes, but gradually increased with rising exposure periods. The blood parametric levels of TLC, ESR and MCHC were increased significantly. The level of blood glucose was increased significantly at $P < 0.01$. It was lower during 15 days but higher in 45 days exposure period.

Metasystox:-

In acute toxicity test the Hb% and TEC were increased significantly after 24, 48, 72 and 96 hours exposure periods at $P < 0.01$ and $P < 0.001$. The values of PCV, MCH and MCV were significantly increased at $P < 0.01$ and $P < 0.001$. TLC, ESR and MCHC were decreased significantly. The biochemical parameters like glucose level also increased in comparison to control fishes. The levels of glucose increased significantly at $P < 0.01$ in treated fishes.


During chronic test the levels of Hb%, TEC, PCV, MCH and MCV were significantly increased at $P < 0.01$ in treated fishes than untreated ones after the end of 15 days, 30 days and 45 days intoxication of metasystox. A significant decreased level of TLC, ESR and MCHC were observed at $P < 0.01$ and $P < 0.001$. The blood glucose level was increased significantly in the insecticidal treated fishes.

Imidacloprid:-

In case of imidacloprid same results were observed as glyphosate and phosphamidon i.e. Hb% TLC, PCV, MCH and MCV decreased significantly while TLC, ESR, MCHC and glucose were increased after 24, 48, 72 and 96 hours exposures.

In chronic toxicity bioassay the Hb % TEC, PCV & MCH decreased significantly after 15 days, 30 days and 45 days exposure of imidacloprid (1/10 of 96 hours LC_{50} concentration). The level of TEC was decreased but no significant difference was observed after the end of 15 and 30 days of exposure periods. A statistically insignificant increased was observed in erythrocytes sedimentation rate except 15 days exposure period which was significant at $P < 0.001$. The level of blood glucose was increased significantly ($P < 0.01$) in treated fishes.

Fishes lives in very intimate contact with their environment. They are susceptible to physical and chemical changes which may be reflected in their blood components. The changes in water quality can alter the haematological and biochemical parameters. Water temperature, O_2 , pH, and other factors also causes disturbance in metabolic rate. Hence this study was also performed to obtain data on seasonal changes in blood parameters of fresh water fishes *Channa punctatus*. If only control fishes were taken into consideration, it was found that Hb%, TEC, TLC, PCV, ESR, MCV, MCH, MCHC and blood glucose level were minimum during summer and maximum during rainy season. When compared to untreated control using Glyphosate, Phosphamidon, Metasystox and Imidacloprid. The activity of Hb%, TEC, PCV, MCH and MCV were decreased while TLC, ESR, MCHC and blood glucose were increased. In case of metasystox treated fishes Hb%, TEC, PCV, MCV and MCH were increased but TLC, ESR and MCHC were decreased. Blood glucose level was also significantly increased. It was found that activity levels of all the parameters were decreased during summer season comparatively.



*All the acute and chronic toxicity experiments were carried out during winter, summer and rainy seasons. Comparisons of different haematological and bio chemical parameters were presented graphically among different season.

BIBLIOGRAPHY

- Abidi R., and U.S., Srivastava (1988) : Effect of endosulfan on certain aspects of haematology of the fish. *Channa punctatus*. Proceedings of National Academy of Science India 58: B (1):55-65.
- Adams S.M., (1990) : Biological indicators of stress in fish. AFS (American Fisheries Society) Symposium 8. American Fisheries Society, Bethesda, MD.
- Agrawal S. J., and A. K., Srivastava (1980) : In effect of endosulfan on certain aspects of haematology of the fish, *Channa punctatus* Toxicology 17:97.
- Agricultural Canada (1991): Pre-harvests use of glyphosate herbicide Discussion document. Pesticides Directorate, Ottawa, Ontario. Amino methylphosphonic acid in water and sediments of boreal forest ponds.
- Aguigwo J.N., (2002) : The toxic effect of cymbush pesticide on growth and survival of African catfish, *Clarias gariepinus* (Burchell 182). J. aquat. sci. 17 (2): 81-84.
- Ahmad F., Ali S. S., Sakoki. A. R., (1995) : Sublethal effect of denitol (fenpropathrin), A systematic pyrethroid on fresh water chinies grass crop, *Ctenopharyngodon idella*. Folia. Biol. (43) 151-159.
- Ajani F., (2008): Harmonal and haematological response of *Clarias garipanus* (Burchell 1822) to ammonia toxicity. African journal of Biotechnology Vol. 7 (19) pp. 3466-3471.
- Akira Kakuno and Jiro Koyama (1994) : Comparative studies on fish blood-II, Seasonal variations of haematological parameters in marine teleostei. Bull. Nati. Res. Inst. Fish. Sci. No. 6, 67-77.
- Alam M. N., and D. N., Sadhu, (2001) : Studies on toxicity of kadeet-36 to *Channa striatus* J. Environ. Pollen., 8: 175-1777.
- Alam M. N., (2000): Toxicity of metacides -50 to a paddy field fish *Channa punctatus* (Bloch) Ind. J. Environ. Ecoplan, 3; 701-703.
- Alam M. N., (2002): Toxicity of kadeet-36 to an air breathing fish, *Clarias batrachus* (L) . Natu. Environ. Tech.1:427-430.
- Alkahem H.F., Z.Ahmed A. S, Al-Akel, Shamus M.J.K., (1998): Toxicity bioassay and changes in haematological parameters of *Oreochromis niloticus* induced by Trichlorfon. Arab. Gulf. J. Scient. Res. 16(3): 581-593.
- Allin C.J. and Wilson R.W. (2000): Effects of pre-acclimation to aluminum on the physiology and swimming behaviour of juvenile

- rainbow trout (*Oncorhynchus mykiss*) during a pulsed exposure. *Aquatic Toxicology*, 51: 213-224.
- Anand Kumar A, Tripathy A.P., Tripathy N. K., (2001): Effect of dimecron on the blood parameters of *Heteropneustes fossilis*. *J Environ Bio*, 22 (4), 297-299.
- Anees M.A., (1988): Haematological abnormalities in a fresh water teleost *Channa punctatus* exposed to sub lethal and chronic levels of three organophosphorus insecticides. *International J. Ecol. and Environ. Sci.*, 4: 53-60.
- Anju Kumari and K., Pandey (1990): Herbicides induced alterations on some haematological parameters of fresh water teleost, *Clarioid batrachus* (Linn.). *Aquatic Environmental and Toxicology*. 50 : 212-218.
- Annune P.A., Hbele S.O. and Oladimeji A.A. (1994): Acute toxicity of cadmium to juveniles of *Clarias gariepinus* (Tuegls) and *Oreochromis niloticus* (Trewavas) *J. Environ. Sci. Health*, A29: 1357-1365.
- Anton F.A. (1993): Degradational behavior of the pesticides glyphosate and diflufenzuron in water. *Bulletin of Environmental Contamination and Toxicology* 51:881-888.
- Arasta T., Bais V.S., and Thakur P., (1996): Effect of nuvan on some biochemical parameters of Indian cat fish, *Mystus vittatus* *J. Environ. Biol.* 17: 167-169.
- Armbrust K.L.(2000) : Pesticide hydroxyl radical rate constants: measurements and estimates of their importance in aquatic environments. *Environ. Toxicol. Chem.* 19, 2175-2180.
- Arunachalam S., Palanichamy S., (1982): Sub lethal effects of carbaryl on surfacing behavior and food utilization in the air-breathing fish, *Macropodus cupanus*. *Physiol. Behav.* 1982; 29: 23-27.
- Asztalos B., Nemcsok J., Benedeczky I., Gabriel R., Szabo A., Refaie O. J., (1990) : The effects of pesticide on some biochemical parameters of carp (*Cyprinus carpio*). *Arch. Environ. Contam. Toxicol.*, 1990; 19: 275-282.
- Ayoola S.O., (2008) : Toxicity of glyphosate herbicide on Nile tilapia (*Oreochromis niloticus*) juvenile , *African Journal of Agricultural Research* Vol. 3 (12), pp. 825-834, December, 2008
- Bala Sashi, Sinha Ratna, Sinha MP, Mehrotra P.N., (1994) : Toxicity of sub lethal concentrations of some heavy metal salts on haematology of *Channa punctatus*. III Erythrocyte counts *J Fresh water Bio*6(2) (1994), 187-190 [10 Ref]

- Balasubramanian S., Ramaswami M., (1991) : Effect of pesticide Sevin on acetyl cholinestrace (AChE) activity in different tissues of *Oreochromis mossambicus* (Peters). *J Ecobio*, 3 (2) (1991), 117-122 [15 Ref].
- Barto B. A., and G. K., Iwama. (1991): Physiological changes in fish from stress in aquaculture with emphasis on the response and effects of corticosteroids. *Annual Review of Fish Diseases* 1:3-26.
- Beauvais S.L., Jones S.B., Brewer S.K., Little E.E., (1880): Physiological measures of neurotoxicity of diazinon and malathion to larval rainbow trout (*Oncorhynchus mykiss*) and their correlation with behavioral measures. *Environ. Toxicol. Chem.* 2000; 19: 1880.
- Begum Ghousia (2009) : Toxicity of dimethoate to aquatic organisms: A Biochemical and Bioaccumulation study *Environmental Biology* 40:38-43.
- Beitinger T. L., (1990) : Behavioral reactions for the assessment of stress in fishes. *Great Lakes Research* 16:495-528.
- Beitinger T.L., Freeman L., (1983) : Behavioral avoidance and selection responses of fishes to chemicals. *Residue Rev.* 1983; 90: 35-55.
- Benerjee G., and Rajendranath (1990) : Hematological changes induced by organophosphorous insecticides in fresh water fish *Clarias batrachus* (linn) . *Trop. Freshwat. Biol.* 2: 197-202.
- Benerji G., and Rajendranath (1999) : Histopathological changes in the liver of *Clarias batrachus* (linn) induced by organophosphorous insecticides. *J. Natu. Cons.* 42: 75-79.
- Bhargava S., Dixit R.S. Rawat M (1999) : BHC and malathion induced changes in TLC and DLC of *Channa striatus*. *Proc Acad Environ Bio*, 8 (1) (1999), 91-94 [16 Ref].
- Bhaskar B.R., and K.S. Rao, (1985) : Some haematological parameters of tarpon, *Megalops cyprinoides* (Broussonet) from Visakhapatham harbour. *Matsy*, 11: 63-69.
- Bhatia H.L., (1972) : Toxicity of copper to fish at different pH levels. *J. Inland fish . Soc. India*: 132-134.
- Bhatia N.P., Sandhu G.S, Johal M.S., (2002) : Endosulfan induced changed in blood chemistry of *Heteropneustes fossilis* pollen, *Res*, 21 (3).
- Bhattacharya Lata (1994) : Effect of DDT exposure on melanophores of the fish, *Oreochromis mossambicus*. *J Env Polln*, 1(2) (1994), 53-54 [6 Ref].
- Birendra Kumar; Banerjee V., (1991): Effects of lethal toxicity of sevin (carbaryl) on the blood parameters in *Clarias batrachus* L.

- Himalayan J Environ Zoo*, 6 (1) , 13-17 [30 Ref].
- Black J. A., Birge, W. J., (1980) : An avoidance response bioassay for aquatic pollutants. Water Resources Research Institute Research Report 123. University of Kentucky, Kentucky.
- Blaxhall P. C., (1972): The haematological and biochemical assessment of the health of fresh water fish. A review of selected literature. *J. Fish. Biol.* 32: 593-604.
- Brain O., R.D., (1967): Insecticides section and metabolism. Acdmec press London pp.125-132.
- Braunbeck T., (1994): Sublethal and chronic effects of pollutants on freshwater fish. *Oxford, UK: Blackwell*, 1994.
- Brewer S.K., Little E. E., DeLonay, A.J., Beauvais, S.B., and Jones, S. B. (1999): The use of automated monitoring to assess behavioral toxicology in fish. Linking behavior and physiology. In *Environmental Toxicology and Risk Assessment* 8th Volume.
- Brewer S.K., Little, E.E., DeLonay, A.J., Beauvais, S.L., Jones, S.B., Ellersieck, M.R. (2001) : Behavioral dysfunctions correlate to altered physiology in rainbow trout (*Oncorhynchus mykiss*) exposed to cholinesterase-inhibiting chemicals. *Arch. Environ. Contam. Toxicol.* 40: 70-76.
- Bridges D. W., Cech, J. J., Pedro, D. N., (1976): Seasonal haematological changes in winter flounder, *Pseudopleuronectes americanus*. *Trans. Am. Fish. Soc.*, 1976; 5: 596-600.
- Brooke L (1993) : Acute and chronic toxicity of nonylphenol to ten species of aquatic organisms. EPA 68 C-0034. U.S. Environmental Protection Agency,
- Brooker M. P., and Edwards, R. W. (1975) : Aquatic herbicides and the control of water weeds. *Water Res.* 9, 1-15.
- Brusle J., Gonzalez I., & Anadon G., In Munshi JSD., & Dutta H.M. (1996) : The structure and function of fish liver. *Fish Morphology*. New York, Science Publishers Inc
- Canli M., (1995) : Effects of mercury, chromium, and nickel on some blood parameters in the carp *Cyprinus carpio*. *Turkish Journal of Zoology*. 19:305-311.
- Canli M., (1996) : Effects of mercury, chromium, and nickel on glycogen reserves and protein levels in tissues of *Cyprinus carpio*. *Turkish Journal of Zoology*. 20:161-168.
- Capkin E, Altinok, I, & Karahan, S., (2006) : Water quality and fish size affect toxicity of endosulfan, an organochlorine pesticide, to rainbow

- trout. *Chemosphere*, 64, 1793-1800.
- Carlisle S.M., and J.T., Trevors (1988) : Glyphosate in the environment. *Water Air Soil Pollut.* 39:409-420.
- Casillas, E. and L.S Smith, (1987) : Effect of stress on blood coagulation and haematology in rainbow trout (*Salmo gairdneri*). *J. Fish Biol.*, 10: 481-491.
- Cech J.J. and D.E. Wohlschlag, (1981) : Seasonal patterns of respiration, gill ventilation and haematological characteristic in the striped mullet (*Mugil cephalus*). *Bull. Mar. Sci.*, 31: 112-119.
- Ceron J.J., Sancho, E., Ferrando M.D., Gutierrez C., Andreu E. (1997) : Changes in carbohydrate metabolism in the eel *Anguilla anguilla*, during short-term exposure to diazinon. *Toxicol. Environ. Chem.* 60: 201-210.
- Chakraporti P., (1986) : Effect of pesticides on the periferial haemogram in an air breathing fish , *Channa punctatus*. Doc. Thesies Patana University. Patana
- Chandra Smita, Ram RN, Singh J., (2001) : Toxic effect of carbofuran on certain hematological parameters in yearlings of *Cyprinus carpio*. *Aquacult*, 2(2) (2001), 137-140 [11 Ref].
- Chandrasekhar and S., Jayabalan (1993) : Haematlogical and biochemical responces of the commen corp *Cyprinus carpio* Linnseus exposed to endosulphsn . University of agriculture science , Bangalore.
- Chandrasekhar S and N., Natrajan (1993) : Endosulphan induced haematological and biochemical alterations in a fresh water cat fish, *Heteropneustes fossilis*. *Ecotoxicology Environment Safety*, 5:329-340.
- Charistoforides C., & Hedley-Whyte, J., (1969) : Effect of temperature and hemoglobin concentration on solubility of O₂ in blood. *J . appl. Physiol.* 27 (5), 592-596.
- Chaturvedi L.D., and Agrawal K., (1993) : Haematological changes in *Heteropneustes fossilis* following exposure to alachlor and rogar. *Advance Biosphere* , 12 (11) 85-92.
- Chaudhary D.K., A.K. Sadhu and P.K., Mukhopadhyay (1984) : Toxicity of two organophosphorous insecticides melathion and Phosphamidon to the fish *Channa striatus*. *Environ. Microbial.* 42: 641-648.
- Chindah A. C., Sikoki F. D. and Vincent Akpu I. (2000) : The effect of organochlorine pesticides on juveniles of a common wetland fish *Tilapia guineensis*. *J. Agric. Biotech. Environ.*, 1 (2): 75- 82.
- Choudhary M.V., Tantarapale V.T., Lokhande P.J., Sherekar P.Y., Kulkarni

- K.M., (1993) : HCH impact on feeding and growth of the freshwater fish *Rasbora daniconius* (Ham). *Uttar Pradesh J Zoo*, 13(2) (1993), 139-142.
- Choudhary N., Sharma M., Verma P., & Joshi S. C. (2003): Hepatopatological and nephrotoxicity in fishes exposed to endosulfan. *J Env Biol*, 24, 305-8
- Claassen M., (1999): "Ecological Risk Assessment as a Framework for Environmental Impact Assessment", *Water Science Technology*. 39(1999), 151-154.
- Cleveland., L, Little, E. E., Ingersoll, C. G., Wiedmeyer, R. H., and Hunn, J. B., (1991) : Sensitivity of brook trout to low pH, low calcium and elevated aluminum concentrations during laboratory pulse exposures. *Aquatic Toxicology* 19:303-317.
- Collazos., M.E., E. Ortega, C. Barriga and A.B. Rodriguez, (1998) : Seasonal variation in hematological parameters male and female *Tinca tinca*. *Mol. and Cell. Biochem.*, 183: 165-168.
- Comes R.D., V.F. Bruns, and A.D. Kelley. (1976a) : . Residues and persistence of glyphosate in irrigation water. *Weed Science* 24(1):47-50.
- Congleton J.L., and La Voie W.J., (2001) : Comparison of blood chemistry values of samples collected from juvenile chinook salmon by three methods. *J Aquat Health*. 13:168-172.
- Connors T.T., Schneider, M.J. Genoway, R.G. and Barraclough, S.A. (1978): Effect of acclimation temperature on plasma levels of glucose and lactate in rainbow trout, *Salmo gairdneri*. *J. Exp. Zool.*, 206: 443- 449.
- Corbett J.R. (1974) : *The Biochemical Mode of Action of Pesticides*. New York: Academic Press, 1974.
- Coutant C.C. (1972): Biological aspects of thermal pollution. II. Scientific basis for water temperature standards at power plants. *CRS Critical Reviews in Environment Control* 3(1) 1:24.
- Cox C., (1998) : Glyphosate (roundup). *J. Pestic Reform*, 18: 3-17.
- Cox L, Koskinen W.C., Yen P.Y. (1997): Sorption-desorption of imidacloprid and its metabolites in soils. *J Agric Food Chem* 45:1468-1472.
- Cranmer M.F., (1986): Carbaryl. A toxicological review and risk analysis. *Neurotoxicology* 1:247-332, 1986.
- Cura J. J., (1998): "Ecological Risk Assessment", *Water Environment Research*. 70(1998), 968-976.

- Dabral R. and L.D., Chaturvedi (1984) : Folidol induced changes in some hematological values of *Heteropneustes fossilis* at low temperatures. J. Adv. Zool., 4 (2) : 92-96.
- Dalela R.C., Bhatnagar, MC, Tyagi, AK, & Verma, SR. (1979) : Histological damage of gills in *Channa gachua* after acute and subacute exposure to endosulfan and rogor. *Mikroskopie*, 35, 301-7
- Dange A.D., (1986): Changes in carbohydrates metabolism in Tilapia, *Oreochromis (Sarotherodon) mossambicus*, during short-term exposure to different type of pollutants. Environ. Pollut., 41(A): 165-177.
- Das B. K., (1998): Studies on the effect of some pesticides and commonly used chemicals on Indian major carps and their ecosystem D. Phil Thesis, Orissa University of Agriculture and Technology, Bhubaneswar (India) 139-162 pp.
- Das B. K., and Mukherjee S. C. (2000) : Sublethal effect of quinalphos on selected blood parameters of *Labeo rohita* (Ham.) fingerlings. *Asian Fish. Sci.*, 13: 225-233.
- Das B. K., and S. C. Mukherjee (2003) : Toxicity of cypermethrin in *Labeo rohita* fingerlings: Biochemical, enzymatic and haematological consequences. *Comp. Biochem. Physiol.*, 134C: 109-121.
- Das P. C., S. Ayyappan and J. K. Jena (2003): Oxygen consumption pattern of fingerlings of Indian major carps, *Catla catla* J Fish Res Bd 29:33-36.
- Das P. C., S. Ayyappan J. K. Jena and B. K. Das, (2004) : Nitrite toxicity in *Cirrhinus mrigala* (Ham.): Acute toxicity and sub-lethal effect on selected haematological parameters. *Aquaculture*, 235: 633-644.
- Das R., (2000) : The toxicity of carbil to aquatic fauna and its impacts on pond ecosystem. Proc. Acad. Environ. Biol. 9: 37.
- Davy F.B., H. Kleerekoper, P. Gensler. (1972) : Effects of exposure to sublethal DDT on the locomotor behavior of the goldfish (*Carassius auratus*). J Fish Res Bd Can 29:1333-1336.
- De La Torre F.R., Salibian A, and Ferrari L. (2000) : Biomarkers assessment in juvenile *Cyprinus carpio* exposed to waterborne cadmium. Environ Pollut. 109:277-282.
- Delor M.E., L. A. Taylor, S.A., Lund P.L., Peniqton, E.D., Strozer and M.H. Fulton. (2000) : Toxicity and biconcentration potential of the agricultural pesticides endosulphan in phytoplankton and zooplankton. Arch. Environ. Contam. Toxicol. 42: 173-181.
- Denton J.E. and M.K., Yousef (1995) : Seasonal changes in hematology of

- rainbow trout, *Salmo gairdneri*. Comp. Biochem. Physiol., 51: 151-153.
- Department of Pesticide Regulation (DPR). (2001b): Case Reports Received by the California Pesticide Illness Surveillance Program, 1982-2001 In Which Health Effects Were Definitely, Probably, or Possibly Attributed to Exposure to Imidacloprid, Alone or In Combination.
- Dethloff G.M., Schlenk D, Khan S, and Bailey HC. (1999) : The effects of copper on blood and biochemical parameters of rainbow trout (*Oncorhynchus mykiss*). Arch Environ Contam Toxicol. 36:415-423.
- Devi M., and Fingerman M. (1995) : Inhibition of acetylcholinesterase activity in the central nervous system of the red swamp crayfish, *Procambarus clarkii*, by mercury, cadmium and lead. Bull. Environ. Contam. Toxicol., 55, 746- 750.
- Dhembare A.J., and Pondha G.M. (2000) : Hematological changes in fish *Punctius sphore* exposed to some fungicides. Ind. J. Environ. Ecoplan. 2: 163-166.
- Dhembare A.J., and Pondha G.M. (2000) : Hematological changes in fish *Punctius sphore* exposed to some insecticide. J. Expt. Zoo, India. 3 (1) : 41-44.
- Dhilon S. S., and Gupta A. K., (1983) : A clinical approach to study the pollutants and intoxication of fresh water fish *Cla*
- Dickman M.D. and K.M. Leung (1998) : Mercury and organochlorine exposure from fish consumption in Hong Kong. Chemosphere, 37: 991-1015.
- Dieter C. D., Duffy, W. G. and Flake L. D. (1996). The effect of phorate on wetland macro invertebrate. Environ. Toxicol. & Chem. 15:(3), 308 - 321.
- Dodson S.I., Hanazazto T. and Gorski P.R. (1995) : Behavioral responses of *Daphnia pulex* exposed to carbaryl and *Chaoborus* kairomone. Environ. Toxicol. Chem. 14 (1): 43-50.
- Doving K.B. (1991) : Assessment of animal behavior as a method to indicate environmental toxicity. comparative biochemistry and physiology. 100C:247-252.
- Dragomieaseu I., Rail L., and Ababai L. (1975) : The effect Carbatox on glycolysis and the activity of some enzymes in carbohydrates metabolism in fresh water fish . Water Res. (9) : 205-209.
- Dubale M.S., and Punita shah (1981) : Biochemical alterations induced by cadmium in the liver on *Channa punctatus*. Environmental research.

- 26 (1):110-118.
- Dutta H.M., I.N., Munshi, J.S.D, & Dutta, HM (Eds.). (1996): A composite approach for evaluation of the effects of pesticides on fish. *Fish Morphology*. New York: Science Publishers Inc
- E.P.A. (1993) : .Glyphosate. R.E.D. Facts. Prevention, Pesticides and Toxic Substances. EPA- 738-F-93-011.
- Ecobichon D.J., I.N., Klaassen C.D., Amdur M.O., & Doull J (Eds) (1996) : Toxic effects of pesticides. *Toxic Agents of Casaretti and Doull's Toxicology: The Basic Science of Poisons* (pp.643-89). New York: McGraw-Hill
- Eddy F. B., (1973): Oxygen dissociation curve of the blood of the tinca, tinca tinca. J.Exp. Biol.58, 281-293.
- Edsall C. C. (1999): A blood chemistry profile for lake trout. J. Aq. Animal Health 11: 81-86.
- Edwards C. A., & Fusher, S. W. (1986) : The use of Cholinesterase measurements in assessing the impacts of pesticides on terrestrial and aquatic invertebrates. In P. Mineau, ed. Cholinesterase – inhibiting insecticides. Their impact on wild and the environment. Elsevier, Amsterdam, The Netherlands, pp. 255 – 275.
- Edwards R., Ferry D.G., Temple W.A. (1996) : 1 *Handbook of Pesticide Toxicology. Classes of Pesticides*. Vol 3. New York: Academic Press, pp 1409–1470.
- Eifac (1983) : European Inland Fisheries Advisory Commission. Revised Report on fish toxicity testing procedures. EIFAC Technical Paper. No. 24
- Ellis A.E., (1981) : Stress and the modulation of defence mechanisms in fish. In: Stress and fish. Pickering AD (Ed). Academic Press London, pp. 147-171.
- Ezzat A.A., M.B. Shabana and A.M. Farghaly, (1993) : Studies on the blood characteristic of *Tilapia zilli* (Gervais). J. Fish Biol., 6: 1-12.
- Feng J.C., D.G. Thompson and P.E. Reynolds. (1990) : Fate of glyphosate in a Canadian forest watershed: 1. Aquatic residues and off-target deposit assessment. Journal of Agricultural Food Chemistry 38:1110-1118.
- Fernando M.D., and E.A. Moliner, (1991) : The effects of infected with *T. brucei*. Rev. Elev. Med. Vet. Pays Trop., time on physiological changes in eel *Anguilla anguilla* 47: 53-57.
- Ferrari Ana , Venturino, Andres, and Pechen de D'Angelo, Ana M. (1989) : Effects of carbaryl and azinphos methyl on juvenile rainbow trout

- Ooncorhynchus mykiss*) detoxifying enzymes.
- Flos R., Tort L, Balacch J (1987): Effects of Zinc Sulphate on haematological parameters in the dogfish (*Scyliorhinus canicula*) and Influence off MS 222. Mar. Environ. Res. 21: 289-298.
- Folin and Wu. (1996) : Determination of blood glucose level. Journal of Biological Chemistry 38:81
- Folmar L. C., H. O. Sanders, and A. M. Julin. (1979) : Toxicity of the herbicide glyphosate and several of its formulations to fish and aquatic invertebrates. Arch. Environ. Contam. Toxicol. 8:269-278.
- Folmar LC, Sanders HO, Julin AM (1979) : Toxicity of the herbicide glyphosate and several of its formulations to fish and aquatic invertebrates. Arch Environ Contam Toxicol 8:269-278. (They tested the toxicity of surfactants separately).
- Forest Service, (1984) : Pesticide Background Statements, Vol. I Herbicides. United States Dept. of Agriculture, Agriculture Handbook No. 633.
- Forstner U., and Prosi F (1989) : Heavy metal pollution in freshwater ecosystem. In "Biological aspects of Freshwater Pollution" : In O. Ravera (ed.), Pergamon Press, Oxford) 129-161.
- Fry F.E.J., (1971) : The effect of environmental factors on physiology of fish. In: Fish Physiology, W. S. Hoar and D. J. Randall (Editors), Vol. 6:1-99, Academic Press, New York.
- Fryday S.N.L., Andrw, D. M., Hart, A. D. M., and Langton S. D. (1996) : Effects of exposure to an organophosphate pesticide on the behavior and use of cover by Captive starlings. *Environ. Toxicol. and Chem.* 15: (9)1590 - 1596.
- Garcia-Santos S, Fontainhas-Fernandes A, and Wilson JM. (2006) : Cadmium tolerance in the Nile tilapia (*Oreochromis niloticus*) following acute exposure: Assessment of some ionoregulatory parameters. *Environ Toxicol.* 21:33-46.
- Gaur K, Pandey Surendra D (1992) : Studies on the toxicity of two organophosphorous compounds on a freshwater teleost, Labeorohitallam.). *Acta Ecologica*, 13 (2) (1992), 82-89 [7 Ref].
- Geol K.A., and Maya (1986): Haematological abnormalities in *Clarius batrachus* (L) Him. *Advance Biosphere*, 5 (11) 187-192.
- Ghosh and Chatterji (1989) : Influence of nuvan on the organic reserve of Indian fresh water fish murrel, *Channa punctatus*. *J. Environ. Biol.* 10: 93-99.
- Ghosh R., Shrotri RV (1992) : Blood glucose and tissue glycogen

- interrelationship in *Scyllaserrata* (Forsk.) chronically exposed to thiodan. *J Environ Bio*, 13 (3) (1992), 233-237 [Ref].
- Ghosh T. K. (1989) : Toxic impact of three organophosphates on carbohydrate metabolism in a freshwater Indian catfish, *Clarias batrachus*, Proceedings of Indian National Science Congress. 135-142pp.
- Ghosh T. K., (1987): Impact of commercial gread hexagon and sumidon on behaviour and some aspects of carbohydrate metabolism in fish *Cyprinus carpio*. Utter Pradesh, Journal of Zoology, 7. 48-62.
- Giesy J.P., Dobson S, Solomon KR, (2000). Ecotoxicological Risk Assessment for Roundup Herbicide. *Rev Env. Contam Toxicol* 167:35-120.
- Gill T.S., and Pant J.C. (1985) : Erythrocytic and leucocytic response to cadmium poisoning in fresh water fish *Puntius conchoni*. *Environ. Res.*, 36, 327-337.
- Gill T.S., Pandey J., Tewari H. (1990): Sublethal effects of an organophosphorus insecticide on certain metabolite levels in a freshwater fish, *Puntius conchoni*. Hamilton. *Pestic. Biochem. Physiol.* 36: 290-299.
- Gill T.S., Pant J.E. (1981): Effect of sublethal concentration of mercury in teleost *Pontius conchoni*. Biochemical and haematological responses. *Indian J. Exp. Biol.* 19: 571.
- Gill T.S., Pant, J.C., & Pant, J.E (1988) : Gill, liver and kidney lesions associated with experimental exposures to carbaryl and dimethoate in the fish (*Puntius conchoni* am.). *Bull Environ Contam Toxicol*, 41, 71-8.
- Goel K. A., Tyagi, S. K., Awasthi, A. K. (1982) : Effect of malathion on some haematological values in *Heteropneustes fossilis*. *Comp. Physiol. Ecol.* 7: 259-261.
- Goel K.A., A. K. Awasthy and S. K. Tyagi. (1982) : Effect of melathion hematological values in *Hreropneustes fossilis*. *Comp. physiol. Ecol.* 7: 259-261.
- Goel K.A., A.K. Awasthi and J.K., Tyagi, (1981) : Comparative haematological studies in some freshwater Indian fishes. 2. Tierphysiology Tierernahrgu. Futtermittelkde. Muzaffarnagar, 46: 202-206.
- Golow A. A. and Godzi T. A. (1994). Acute toxicity of Deltamethrin and Dieldrin to *O. niloticus*. *Bull. Environ. Contam. Toxicol.* 52: 351-354.

- Gray R.H., (1990) : Fish behavior and environmental assessment. *Environmental Toxicology and Chemistry* 9:53-68.
- Grillitsch B; Vogl, C. & Wytek, R. (1999) : Qualification of spontaneous undirected loco motor behavior of fish for sublethal toxicity testing. Part II. Variability of measurement parameters under toxicant induced stress. *Environ. Toxicol. & Chem.* , 18(12): 2743 – 2750.
- Gross R., Gliwitzki M., Gross P. and Klaus F. (1996) : Food and Nutrition Bulletin Volume 17, Number 1, United Nations University Press.
- Gubbins M.J., Eddy, F.B., Gallacher, S. and Stagg, R.M. (2000): Paralytic shellfish poisoning Toxins induce xenobiotic metabolism enzyme in salmon (*Salmo salar*). *Marine environmental research* 50: 469-483.
- Gupta A. K. and A. Amuni., A. Ramjana and R.C. Dalela (1995) : Toxic effect of clorodence and melathion on certain haematological parameters of a fresh water teleost , *Notopterus notopterus* J. *Environ. Biol.* 16: 219-223.
- Gupta D. K., Rai U.N., Singh A. and Inouhe M. (2003) : Cadmium accumulation and toxicity in *Cicer arietinum* L. *Poll. Res.*, 22, 457-463.
- Guzelian P. S., (1982) : Comparative toxicology of chlordecone (kepone) in humans and experimental animals. *Annu Rev Pharmacol Toxicol* 22:89-113, 1982.
- Haley T. J., (1979) : Review of the toxicology of paraquat (1,19-dimethyl-4,49-bipyridinium chloride). *Clin Toxicol* 14:1-46, 1979.
- Harding J., and Hogland, L. B., (1984): "Seasonal variation in blood components of reared Baltic salmon (*Salmo salar*)" *J.Fish.Biol.*, 24:565-579.
- Hart W.B., Doudoroff, P. and Greenbank J (1985) : The evaluation of the toxicity of industrial wastes, chemicals and other substances of freshwater fish. Atlant Refining Co., Phil. 317 pp.
- Heath A. G. (1995) : Water Pollution and Fish Physiology, 2nd ed. CRC, Boca Raton, FL, USA.
- Heath A.G. (1996) : Water pollution and fish physiology. CRC. Press. Inc. Boca Raton, Florida,. 245p.
- Hildebrand L. D., D. S. Sullivan, and T. P. Sullivan. (1980) : Effects of RoundupR herbicide on populations of *Daphnia magna* in a forest pond. *Bull. Environ. Contam. Toxicol.* 25:353- 357.
- Hinton D.E. and Lauren D.J. (1990) : Integrative histopathological effects of environmental stressors on fishes. *Amer Fish Soc Symp*, 8, 51-66.

- Hoar S. K. , Blair A., Holmes F. F. (1986) : Agricultural herbicide use and risk of lymphoma and soft-tissue sarcoma. *JAMA* 256:1141-1147, 1986.
- Hochachka P.W., Guppy, M., Guderly, H.E., Storey K.B., and Hulbert W.C. (1978) : Metabolic biochemistry of water versus air breathing fishes muscle enzymes and ultrastructure. *Can. J. Zool.*, 56: 736-750.
- Hoemechaudhuri S., Pandit T., Podder S., and Benerjee S. (1986) : Effect of mahuva oil cake on the blood cells and blood values of an air breathing cat fish *Heteropneustes fossilis* and a carp *Cyprinus carpio* *Proc. Indian Acad. Sci.* 95; 617-622.
- Hoff P.T., Van Dongen W., Esmans E.L., Blust R. and De Coen W.M. (2003) : Evaluation of the toxicological effects of perfluorooctane sulfonic acid in the common carp (*Cyprinus carpio*). *Aquatic Toxicology* 62:349-359.
- Howerton R. (2001): Best Management practises For Hawaiian Aquaculture. Center for Tropical and Subtropical Aquaculture, Publication 148; 7-31.
- Hrudey S.E., W., Chen and C.G. Roussex, (1996): Bioavailability in environmental risk assessment. CRC Press, Inc, Lewis publishers, USA. pp: 125- 133.
- Hughes G.M., (1981) : Effects of low oxygen and pollution on the respiratory systems of fish. (AD Pickering ed). Academic Press, London, pp. 121-146.
- Imbamani N., and R, Shrivassan (1998): Effect of phosphamidon toxicity and pesticidal histopathology of the fish *Sarotherodon mossambica*. *J. Ecotoxicol. Environ. Monit.* 8: 85-95.
- Ipinmoroti K.P., A.A. Oshodi and R.A. Owolabi, (1997): Comparative studies of metals in fish organs, sediment and water from Nigerian fresh water fish ponds. *Pak. J. Sci. Ind. Res.*, 40: 70-74.
- Iwama G. K., A. D. Pickering, J. P. Sumpter, and C. B. Schreck (1997) : Fish stress and health in aquaculture. society for experimental biology, Seminar Series 62. Cambridge University Press, Cambridge, U.K.
- Jagdish Mishra and Anil K. Shrivastava (1981): Melathion induced haematological and biochemical changes in indian cat fish *Heteropneustes fossilis*. *Environmental research* 30,393-3.98.
- Jai Nath Rao., Ch Madhu., V. Ananda Rao V. S. R. Murthy (1984): Phosphamidon toxicity to fresh water fish, *Sarotherodon mossambicus* . *J. Environ, Biol.*, 5 (3), 157-163.

- Jain R., and K.D., Mishra. 1993: Acute toxicity of butachlor to *Puntius titico* a teleost. *Bull. Environ. Sci.* 11: 1-5.
- Jain R., and K.D., Mishra. 1995: Acute LC₅₀ of 2, 4-dichlorophenoxyacetic acid to fish *Rasbora deniconius* comparative evolution by static bioassay and residual oxygen bioassay test. *J. Environ Biol.* 16: 105-111.
- Janardhana Reddy., S., Kalarani, V., Tharakanadha B., Reddy, D. C., Ramamurthi R., (1978): Changes in energy metabolism of the fish, *Labeo rohita* in relation to prolonged lead exposure and recovery. *J. Ecotoxicol. Environ. Monit.*, 1998, 8 (1), 43-53.
- Jebakumar S.R.D. , A.K. Kumar guru and J. Jayabalan (1997) : Accumulation and dissipation on phosphamidon in tissues of fresh water fishes *Oreochromis mossambicus*. *Comp. Physiol. Ecol.* 18: 12-17.
- Jeyaratnam J., De Alwis Seneviratne R. S, Copplesstone J. F., (1982) : Survey of pesticide poisoning in Sri Lanka. *Bull WHO* 60:615-619, 1982.
- Jhon P.J., Rathor. And A. Prakash (1989) : Metasystox generated hematological anomalies in *Heteropneustes fossilis*. *Nat. Dev. Natu. Resour. Conserv.* 160-165.
- Jhosi P. and Deep H. (2002) : Effect of linden and malathion exposure to certain blood parameters in a fresh water teleost fish *Clariatus batrachus*. *Poll. Res.* (21) 55-57.
- Joshi P.C., (1989) : Seasonal changes in the blood parameters of a hillstream teleost, *Channa gachua*. *Comp. Physiol. Ecol.*, 14: 71-73.
- Josif Jhon P. (2007) : Alteration in certain blood parameters of fresh water teleost *Mystus vittatus* after chronic exposure to metasystox and sevin. *Fish physiol Biochem* (2007) 33: 15-20.
- Jyothi B. and Natrajan G. M., (1999) : Certain pesticides induced carbohydrate metabolic disorder in the serum of fresh water fish *Clarias batrachus* (Linn). *Food and Chemical toxicology* 37, 417-421.
- Kamble G.B. and Muley D.V. (2000) : Effect of acute exposure of endosulfan and chlorpyrifos on the biochemical composition of the fresh water fish *Sarotherodon mossambicus*. *Indian J Environ Sci*, 4(1) (2000), 97-102 [19 Ref].
- Kaplan H.M., and Crouse G.T. (1956) : Blood change underlying the seasonal resistance to frogs to disease. *Capio*, pp. 52-56.
- Kashiwada S., Ishikawa, H., Miyamoto, N., Ohnishi, Y. and Magara, Y.,

- (2002) : Fish test for endocrinedisruption and estimation of water quality of Japanese rivers. *Water Res.*, 36: 2161-2166.
- Kemann H., (1981) : Fish toxicity test with mixtures of more than two chemicals: a proposal for a quantitative approach and experimental results. *Toxicology* 19: 229-238.
- Khalid A., Al-Ghanim, Hmoud F. Al-Khem Al-Balawi, Ali S. Al-Akel, Fahad Al- Mansid , Zubair Ahmad (2008) : Ethological response, hematological and biochemical profile of carp (*Cyprinus carpio*) exposed to triclofon. *Jouranal of food, Agricultural and Environmenta*; Vol., 6 (3 & 4) 173-179.2008.
- Khangarot B.S. , Sehgal and M.K., Bhasin (1985) : Effect of pH on toxicity of sodium pentacholorophenate to fry of common carp in soft water. *Arch. Hydrobiol.* 103(3) : 375-379.
- Khangarot B.S., and P.K. Ray (1988e) : Study on the effect of malathion to fresh water teleost, (Bloch) *Channa striatus* and *Puntius sophoer* (Hamilton). *Arch. Hydrobiol.* 470: 1-5.
- Khangarot B.S., and P.K., Ray (1988b) : Sensitivity of fresh water pulmonate snails, *Lymnaea leutiola* L, to heavy metals. *Bull. Environ. Contam. Toxicol.* 41 : 208-213.
- Khangarot B.S., R. Takkroo, R.R., Singh and S.P. Shrivastava. (1991) : Toxicity of biconcentration of Hexachloroxen (HCH) in an air breathing catfish , *Secobranchus fossilis* (Bloch) .*Bull. Environ. Contam.* 47: 904-911.
- Khattak I. U. D., Hafeez, M. A. (1996) : Effect of malathion on blood parameters of the fish, *Cyprinion watsoni*. *Pak. J. Zool.* 28: 45-49.
- Kidd H., James DR (1991) : The Agrochemicals Handbook, 3rd ed. The Royal Society of Chemistry, Unwin Brothers Limited, Old Woking, Surrey, UK Chem 16:336-345.
- Kleerekoper, H., Waxman J.B., and Matis J. (1973) : Interaction of temperature and copper ions as orienting stimuli in the locomotor behavior of the goldfish (*Carassius auratus*). *Journal of the Fisheries Research Board of Canada* 30:725-728.
- Kori Siakpers Ovei, Adamu Kabir Mohammed and Madukelum Ifeanyi Theresa (2007) : Acute haematology effects of sublethal levels of paraquate on the African cat fish *Clarias garipinus*. *Research journal of Environmental Science* 1 (6) ; 331-335.
- Kori-Siakpere O, Egor VE (1999). Haematological characteristics of African mudfish. *Clarias buthupogon*. (Pisces: Clariidae). *Bull. Sci. Assoc. Niger.* 21: 177-185.

- Kori-Siakpere O., (1985): Haematological characteristics of *Clariaies sheriensis* (Sydenham). J. Fish Biol. 27: 259-263.
- Kori-Siakpere O., (1991): Chronic sublethal haematological effects of copper in fresh water teleost. *Clarias isheriensis* (Sydenham). Niger. Ann. Nat. Sci. 1: 37-43.
- Koyama J., Ozaki H (1984) : Haematological changes of fish exposed to low concentrations of cadmium in the water. Bull. Jpn. Soc. Sci. Fish. 50: 199-203.
- Kulshrestha S.K. and Arora, A., (1984) : Impairment caused by sublethal doses of two pesticides in ovaries of freshwater teleost *Channa striatus* Publ-Dept. Zool. Toxicol 20(1): 93-98.
- Kumar B., and Banerji V., (1990): Effect of sublethal toxicity of sevin on blood parameters in *Clariious batrachus* (L) Him. Journal of environmental Zoology, (4) 166-172.
- Kumar B.K., (1991): Histopathological and hematological changes induced by pesticides on fresh water teleost fish, *Clarias bacrachus* doctoral thesis, Patna univ. Patana, India.
- Kumar G.S., and D.J. Nelson.(1997) : Effect of Ekalux on haematology of the freshwater fish *Etroplus maculatus*. Journal of Ecotoxicological and Environmental Monitoring 7:113-116.
- Kumar Hemant and Gupta A.B., (1997): Toxicity of organophosphorus corbanate and synthetic pyrethroid pesticides to the Indian cat fish (*Heteropneustes fossilis*) J. Nature, conserve.9. (1), 111 - 114.
- Kumar S., Lata S., and K., Gopal(1999): Deltramethrin induced physiological changes in fresh water Cat fish .Bull. Environ. Contam. Toxicol. 62: 254-258.
- Kumar Suresh, Lata Swarn, Gopal Krishna (1999) : Acute toxicity of deltamethrin to the freshwater teleosts *Heteropneustes fossilis* and *Channa punctatus*. Proc Acad Environ Bio, 8(1) (1999), 83-85 [14 Ref].
- Kumar V.A., Karenmungikar and S. L. N. Reddy., (1997): A comparative study of sub lethal effects of endosulphan and cypermethrin on blood serum profiles in air breathing cat fish *Clarias batrachus* (Linn.) J. Environ. Polln. 4: 131-135.
- Kuruppasamy R., Subathara S., and Puvaneswari S. (2005): Haematological responses to exposure to sub-lethal concentration of cadmium in air breathing fish, *Channa punctatus* (Bloch). J. Environ. Biol., 26(1), 123-128.
- Kutty M.N., Raj S.P., and Jeyachandran P. (1977) : Comparative tolerance

- of freshwater fishes to selected pesticides. In: Seminar on Environmental Impact on Development and Adoption of Suitable Guidelines for Project Control in Selected Sectors. Inst. Engin. Hyderabad. (India).
- Lakota S., Raszka, A and Roszkowski J., (1983) : D.D.T., Lindane and Toxaphene exerted toxic effects on carp (*Crprinus carpio*) fry at concentrations 1-10 ppm exposed for 24 hours, Folia Biology. 31 (1):93-99.
- Lal H., Misra V., Viswanathan, P.N., and Krishnamurthi C.R., (1984): Effect of synthetic detergents on some of the behavioral patterns of fish fingerlings, *Cirrhinus mrigala* and its relations to ecotoxicology, Bull. Environ, Contam, Toxicol 32, 109-115.
- Lata S., K. Gopal and N.N. Singh. (2001) : Toxicological and morphological studies in a cat fish, exposed to carbyle and carbofuran. J. Ecophysiol. Occup.Hilth. 1: 109-120.
- Lavin S., Cuenca, R., Marco, I., Velarde, R. and Viñas, L. (1992) : Hematology and Blood chemistry of the Marsh Harrier (*Circus aruginosus*). Comparative Biochemistry and Physiology 103A (3):493-495-72.
- Little E.E., and S. E., Finger. (1990) : Swimming behavior as an indicator of sublethal toxicity in fish. Environmental Toxicology and Chemistry 9:13-19.
- Little, E.E., Dwyer, F. J., Fairchild, J.F., DeLonay, A. J., and Zajicek, J.L. (1993b) : Survival and behavioral response of bluegill during continuous and pulsed exposures to the pyrethroid insecticide ES-fenvalerate. Environmental Toxicology and Chemistry 12:871-878.
- Liu M., Lanford, J. and Casida, J. (1993) : Relevance of [³H]imidacloprid binding site in house fly head acetylcholine receptor to insecticidal activity of 2-nitromethylene and 2-nitroiminoimidazolidines. Pestic. Biochem Physiol. 46:200-206.
- Liu M-Y, Casida J. E., (1993): High affinity binding of [³H] imidacloprid in the insect acetylcholine receptor. *Pestic Biochem Physiol* 46:40-46, 1993.
- Liu M-Y, Lanford J, Casida (1993): Relevance of [³H] imidacloprid binding site in house fly head acetylcholine receptor to insecticidal activity of 2-nitromethylene and 2-nitroimino-imidazolines. *Pestic Biochem Physiol* 46:200-206, 1993.
- Lone K. P., and Javed M.Y. (1976) : The effect of sublethal doses of DDT and Dieldrin on the blood of *Channa punctatus*. (Bloch). Pakistan. J. Zool. 8: 143- 149.

- Lone K.P., and M. Y. Javed (1976); The effect of sublethal doses of three organophosphorus insecticides on haematology of *Channa punctatus* (Bloch) Pak. J. Zool., (8) 77-84.
- Lund-Hoie, K, and H. O. Friestad. (1986): Photodegradation of the herbicide glyphosate in water. Bull. Environ. Contam. Toxicol. 36:723-729.
- Luskova V., (1997) : Annual cycles and normal values of hematological parameters in fishes. Acta Sc. Nat. Brno 31(5) : 70-78.
- Luskova V., Savoboda M., Kolaropova J., (2002) : The effect of diazonium on blood plasma biochemistry in carp (*Cyprinus carpio*). Acta. Vet. Brno.71. 117-123.
- Mahajan C. L., and Juneja S., (1979) : Effect of aldrin on peripheral blood of fish (*Channa punctatus*). Ind. J. Environ. Hith. 21. (2) 162-172.
- Mahesh Tyagi, V.P. Agrawal and Nandini (1989): Hematological abnormalities induced in *Ophiocephalus punctatus* by Endosulfan. Environmental series Vol II, 81-84
- Mala F. A., G. Sharma, Singh S. (2009): Chlorpyrios pesticides toxicity on erythrocyte sedimentation rate in fish, *Channa punctatus* (Bloch) Biology of Medicine, Vol 1 (2): 45-55.
- Malik J., G. Barry and G. Kishore. (1989): A mini-review of "The herbicide glyphosate." BioFactors 2(1):17-25.
- Maria Santos Reis Bonorino Figueiredo, Alex J Anderson. (2009) : Digestive enzyme spectra in crustacean decapods (Paleomonidae, Portunidae and Penaeidae) feeding in the natural habitat. *Aquaculture Research* 40:3, 282-291 Online publication date: 1-Mar-2009.
- Marrs R.H., A. J. Frost, R. A. Plant, and P. Lunnis. 1993. Determination of buffer zones to protect seedlings of non-target plants from the effects of glyphosate spray drift. *Agriculture, Ecosystems and Environment* 45:283-293.
- Martinez F.J., M.P., Garcia-Riera, M. Canteras, J. De Costa and S. Zamora, (1994) : Blood parameters in rainbow trout (*Oncorhynchus mykiss*): Simltaneous influence of various factors. Comp. Biochem. Physiol., 101: 95-100.
- Mason C.F., (1996) : Biology of freshwater pollution, 3rd edn., Longman, U.K. pp1-4.
- Mastumura F., Bouch G.M. and Mistol (1972) : Environmental toxicology of pesticides. Academic Press, New York, London. (3) 498-512.
- Mathur S., (1994) : Molluscicide induced toxicity and gonad damage in a

- freshwater pulmonate *Lymnaea luteola* (Lammarek). *Polln Res.* 13(4) (1994), 403-409 [9 ref].
- Matsuda K., Buckingham S.D., Kleijer D., Rauh, J.R., Grauso M. and Sattelle D.B. (2001) : Neonicotinoids: insecticides acting on insect nicotinic acetylcholine receptors. *Trends Pharmacol. Sci.* 22, 573-580.
- Matsuda K., Buckingham, S. D., Freeman, J.C., Squire, M. D., Baylis, H.A. and Sattelle, D. B.(1998) : Effects of the α .subunit on imidacloprid sensitivity of recombinant nicotinic acetylcholine receptors. *Br. J. Pharmacol.* 123:518-524.
- Matsumura F., (1985) : *Toxicology of Insecticides*. New York: Plenum Press, 1985, pp 122-128.
- Matthiessen P., (1981) : Haematological changes in fish following spraying with endosulfan for tsetse fly control in Botswana. *J. Fish. Biol.* 18(4): 461-469.
- Maule A.G., Schreck C.B., (1990): Changes in numbers of leucocytes in immune organs of juvenile coho salmon (*Oncorhynchus kisutch*) after acute stress or cortisol treatment. *J. Aquat. Anim. Health* 2: 298-304.
- Mazeaud M.M. and I. Mazeaud (1981) : Adrenergic response to stress in fish In *Stress and Fish*. Pickering A.D. Eds. Academic Press. New York. Chap.3:150 pp
- Mbiapo F., Youovop G., (1993): Regulation of pesticides in Cameroon. *J Toxicol Environ Health* 39:1-10, 1993.
- Mc Donald M.D. and Grosell M. (2006): Maintaining osmotic balance with an aglomerular kidney. *Comp Biochem Physiol C*. 143:447-458.
- Mc Kim J. M. and Lien G. J. (2001): Toxic responses of the skin in target organ toxicity in marine and freshwater teleosts organs. Eds., Schlenk
- Mc. Donald D.G., Hobe H. and Wood C. M., (1980): The influence of cadmium on the physiological response of the rainbow trout, *Salmo gairdneri*, to low environment pH . *J. Expl. Biol.* 88, 109-131.
- Medda C., Sarkar S.K., Ganguly S., Bhattacharya B., (2002) : Effect of organophosphorus on three important indicators and their recovery in two fresh water carps, *Lebio rohita* and *Cirrhinus mrigala*. *Poll. Res.* 12 (3) ; 151-157.
- Mikko Nikinmaa and Virpi Tervonen (2003) : Regulation of blood haemoglobin concentration Hypoxic fish. *Fish physiology and water quality* EPA/600/R-04/049, 197-206.

- Milligan C.I., and Wood CM., (1982) : Disturbances in haematology , fluid volume distribution and circulatory function associated with the low environment pH in rainbow trout, *Salmo gairdneri*, . J. Expl. Biol. 99, 397-415.
- Minaxi Das and Shambhu Prasad (1996) : Effect of lethal concentration of metasytox (Oxydemeton-methyl) on blood parameters of an air breathing fish *Heteropneustes fossilis* (Bloch) Aquatic Environment and Toxicology. Vol. (32) : 115-121.
- Mishra B. K., (1993) : Effect of some pesticides on different blood parameters of *Clarius batrachus*. Doc. Theses Patana University. Patana
- Mishra C.K., Hakim A, Kumar J (1994) : Effect of monocil pesticides on the oxygen consumption and related blood parameters in *Channa punctatus* (Bloch). *J Freshwater Bio*, 6(1) (1994), 93-99 [27 Ref]
- Mishra D. S., Shrivastava, S.K., Shrivastava and A.K. Shrivastava (2001) : Plasma calcium and inorganic phosphate levels of fresh water cat fish *Heteropneustes fossilis* in response to cypermethrin treatment . J. Ecophysiol. Occup. Health, 1: 131-138.
- Mishra Dilip K, Tripathy Praful C, Hota Ashok K ., (1991) : Toxicity of kilex carbaryl to a freshwater teleost *Channa punctatus* (Bloch). *J Appl Zoo Res*, 2 (2) (1991), 96-98 [10 Ref].
- Mishra J., Shrivastava A. K., (1983) : Malathion induced hematological and biochemical changes in the Indian catfish *Heteropneustes fossilis*. Environ. Res. 30: 393-398.
- Mishra S., Shrivastava AK (1979) : Haematology as an index of sublethal toxicity of zinc in fresh water teleost. Bull. Environ. Contam. Toxicol. 22: 675-698.
- Mishra S., Shrivastava AK (1980) : The effects of copper on the blood of a teleost. Ecotoxicol. Environ. Safety 4: 191-194.
- Mishra U.K., (1988) : Effect of herbicides on major carp seed of Indian major carp, *Catla catla*. Ph.D. Thesis, Bhopal University, Bhopal.
- Mishra U.K., O.N., Seth and K.D., Mishra., (1989) : Toxicity of lasso (alachlor) to the seed of Indian major carp, *Catla catla* Environ. Ser. II : 91-99.
- Mishra U.K., R. Jain, S. Singh , K. Sukla and K. D. ,Mishra (1998) : Toxicity of herbicides to fish; A brief review . pp.252. Ecology of polluted water and Toxicology. Technoscience Publication, Jaipur.
- Mitchell D.G., Chapman P.M., Long T.J., (1987). Acute toxicity of Roundup and Rodeo herbicides to rainbow trout, Chinook, and Coho

- salmon. Bull. Environ Contam Toxicol. 39:1028-1035.
- Mitchell D.G., Chapman P.M., Long T.J., (1987) : Acute toxicity of Roundup and Rodeo herbicides to rainbow trout, chinook and coho salmon. Bull Environ Contam Toxicol 39: 1028-103.
- Mitchell DG, Chapman PM, Long T.L. (1987) : Acute toxicity of round up and Rodeo herbicides to rainbow trout, Chinook and *Coho salmon*. Bull. Environ. Contam. Toxicol. 39:1028-1035.
- Monsanto and Company (1981) (also called Schroeder CA, 1981): A Three-generation reproduction study with glyphosate in rats. MRID No. 00093879. Available from EPA.
- Monsanto Company (1985) : *Toxicology of Glyphosate and Roundup Herbicide*. St Louis, 1985.
- Monsanto Company, (1985): Toxicology of Glyphosate and Roundup Herbicide, Department of Medicine and Environmental Health, St. Louis, MO.
- Morrison M.L. and E.C. Meslow. (1984): Effects of the herbicide glyphosate on bird community structure, western Oregon. Forest Science 30(1):95-106.
- Moses M., (1989) : Glyphosate herbicide toxicity. *JAMA* 261:2549, 1989.
- Muhammad Atamanalp and Telat Yanik (2002) : Alterations in the hematological parameters rainbow trout (*Oncorhynchus mykiss*). Turk. J. Vet. Anim. Sci. 26: 1157-1160.
- Mukherjee M. K., and S.K. Khonr (1984) : Toxicity of copper, zinc and iron to fish, plankton and worms. *Gobes*.11: 204-207.
- Mukhopadhyay P. K., and Ghosh, S., (2000) : Fish in human health and nutrition. *Fishing Chimes*, 20(2): 31-33.
- Mukhopadhyay P. K., and P. V. Dehadrai (1980) : Biochemical changes in the air breathing catfish, *Clarias batrachus* (L) exposed to malathion. *Journal of Environmental Biology* 5:221-229.
- Mukhopadhyay P. K. and Kaushik, S. J., (2001): Nutritional requirements of Indian major carps. *International Aqua. Feed*, January, 2001. 28-32.
- Mukhopadhyay P.K., and, Dehadrai, P.V. (1980): Studies on air breathing cat fish *Clarias batrachus* (Linn) under sublethal exposure. *Ind. J. Exp. Biol.* 18: 348-352.
- Musa S.O., Omoregie E., (1999): Haematological changes in the mudfish, *Clarias gariepinus* (Burchell) exposed to malachite green. *J. Aquat. Sci.* 14: 37-42.

- Nagata K., Aistrup, G. L. Song, J. H. and Narahashi, T. (1996): Subconductance-state currents generated by imidacloprid at the nicotinic acetylcholine receptor in PC 12 cells. *Neuroreport*. 7:1025-8.
- Nagata, K., Song, J. H., Shono, T. and Narahashi, T. (1998) : Modulation of the neuronal nicotinic acetylcholine receptor-channel by the nitromethylene heterocycle imidacloprid. *J. Pharmacol. Exp. Ther.* 285:731-8.
- Naqvi S.M., and R. Howkin, (1988) : Toxicity of selected insecticides (Thiodon, Security, Sportan and Sevin) to mosquito fish, (*Gambusia affinis*). *Bull. Environ. Contam. Toxicol.* 40: 779-782.
- Natarajan G. M., (1989): Changes in the carbohydrate metabolites during acute and chronic exposure of airbreathing fish *Channa striatus* to oxydemeton methyl (metasystox). *Comp. Physiol. Ecol.* 14: 181-184
- Natarajan G.M., (1981): Changes in the bimodal gas exchange and some blood parameters in the air-breathing fish, *Channa striatus* (Bleeker) following lethal (LC50-48h) exposure to metasystox (demeton). *Curr. Sci.* 1981; 50: 40-41.
- Natarajan G.M., (1983) : Effect of sublethal (LC50-48h) concentration of metasystox on selected oxidative enzymes, tissue respiration and histology of gills of the freshwater air-breathing fish, *Channa striatus* (Bleeker). *Curr. Sci.* 1981; 50: 985- 989.
- Nath R., (2003) : Effect of cadmium and zinc on the haematology of an air breathing fish *Clariaus batrachus* (L). *J. Adv. Zool.*, 24(1&2), 57-60.
- Nath. Ravindra, Banerajee V. (1999) : Influence of leathal and subleathal toxicity of Rogor on the blood parameter of fresh water fish *Heteropneustes fossilis*. *Env. Eco.* 17(4) : 940-944.
- National Library of Medicine, (1992): Hazardous Substances Databank. TOXNET, MedlarsManagement Section, Bethesda, MD. Pesticide Fact Sheet: Glyphosate. Office of Pesticide Programs, U.S. Environmental Protection Agency, Washington, DC. EPA Publication No. 540/FS-88-124, 1986.
- Natarajan G. M., (1983) : Effect of lethal concentration of metasystox on hematology and free amino acid and glutamate dehydrogenase in air breathing fish (*Channa punctatus*), *Comp. Physiol. Ecol.* Vol. 8. pp. 254-256.
- Nauen R., Ebbinghaus-Kintscher, U. and Schmuck, R. (2001a.) : Toxicity and Nicotinic Acetylcholine Receptor Interaction of Imidacloprid and its metabolites in *Apis Mellifera* (Hymenoptera: Apidae). *Pest. Manag. Sci.* 57:577-586,

- Nemcsok and Bones (1982) : Comparative study on the sensitivity of different fish species to metal pollution, *Acta . Biol . Hung.* 33: 23-27.
- Neskovic N., Elezovic I, Karan V., Poleksic V. and Budimir M. (1993) : Acute and subacute toxicity of atrazine to carp (*Cyprinus caprio* L). *Ecotoxicol Environ Saf* 25: 173-182
- Neskovic N.K. (1996): Biochemical and histopathological effects of glyphosate on carp, *Cyprinus carpio*. *Bulletin of Environmental Contamination and Toxicology* 56:295-302.
- Neskovic N.K., Poleksic V, Elezovic I, Karan V, Budimir M., (1996) : Biochemical and histopathological effects of glyphosate on Carp. (*Cyprinus carpio* L.). *Bull. Environ. Contam. Toxicol.*, 56: 295-302.
- Neumosok J.G, Hughes G. M., (1998) : The effect of Copper Sulphate on some biochemical parameters rainbow trout. *Environ. Pollut.* 49: 77-85.
- Newton M. (1984) : Fate of glyphosate in an Oregon forest ecosystem. 32:1144-1151.
- Nimmo D.R., Coppage D.L., Pickering Q.H., and Hasan D.J., (1987) : Assessing the toxicity of pesticides to aquatic organisms, in *slint spring Revisited* G.J., Macro R.M., Hollingworth and w., Dhuram, Eds. *American Chemical Society, Washington, DC.* 47-70.
- Nomura N. S., and H. W. Hilton. (1977) : The adsorption and degradation of glyphosate in five Hawaiian sugarcane soils. *Weed Research* 17:113-121.
- Noori M., Cakmak and Asiye Girgn (2003): Toxic effect of synthetic pyrethroid insecticides (cypermethrin) on blood cell of Rainbow trout (*Oncorhynchus mykiss*). *Biological science* 3(8) ; 694-698.
- Nowak B., (1992): Histological changes in gills induced by residues of endosulfan. *Aquat Toxicol*, 23, 65-83.
- Nussey G., (1994): The effect of copper on the blood coagulation and general haematology of *Oreochromis mossambicus* (Cichlidae). *Doctral Thesis, RAU.*
- O'Neal C.C. and Weirich, C.R., (2001) : Effects of low level salinity on prod. and haematological parameters of ,channel catfish, *Ictalurus punctatus* reared in multicrop
- Odieta W.O., (1999) : Environmental physiology of animals and oxygen consumption of *Channa puntatus* (Bloch) and Pollutions. Published by Diversified Resources Proc. Symp. *Environ. Biol.*, pp: 343-348.
- Olla B.L., Pearson W.H. and Studholme A.L. (1980) : Applicantly of

- behavioral measures in environmental stress assessment. Rapp P-V Renn. Cons. Int. Explor. Mer., 179, 162-179.
- Omoniyi I, Agbon A, Sodunk, SA (2002) : Effects of lethal and sub-lethal concentrations of tobacco (*Nicotiana tobaccum*), leaf dust extractum on weight and haematological changes in *Clarias gariepinus* (Buchell 1822). J. Appl. Environ. Man., 6: 37-41.
- Omoriegie E, Ufodike EBC, Keke Í. R., (1990): Tissue chemistry of *Oreochromis niloticus* exposed to sublethal concentrations of gammalin 20 and actellic 25EC. J. Aquat. Sci. 5: 33-36.
- Onusiriuka BC, Ufodike EBC (1998) : Growth of African catfish, *Clarias gariepinus* (Teugels) subjected to sublethal concentrations of water extracts of akee apple, *Blighia sapida* and sausage plant *Kigelia africana*. J. Aquat. Sci. 13: 59-62.
- PAN Pesticide Database (1996) : Chemical toxicology studies on aquatic organisms www.pesticideinfo.org.
- Pandey A.K., Pandey G.C. (2001) : Thiram and Ziram fungicides and induced alterations on some hematological parameters of fresh water cat fish, *Heteropneustes fossilis*. Indian. J. Environ. E. Ecoplan. 5(3):437-442.
- Pandey B.N., S.S. Prasad A.K. Chanchal and S.B. Singh (1984) : Effect of some pesticides on the oxidative metabolism and hematology of *Clarias batrachus* (Linn) Prac. Sen. Eff. Pest. Aq. Face. 13-18.
- Pandey P. K., N. K. Singh and B. P Chaudhary (1980) : Effect of organophosphorous insecticide, malathion on the haematology of *Channa punctatus*. Proceedings of 6th. Indian Science Congress Part II. 213 pp.
- Pandey Reeta, Bhargava S, Saraf DK (1993) : Toxic effect of some biocides on total serum protein in *Heteropneustes fossilis*. Indian J Env Toxicol, 3(1&2) (1993), 5-6 [8 Ref].
- Parish P.R., (1985) : Acute toxicity Test. In: Fundamental of Fish Aqua. Toxicol., 19: 167-204.
- Pascoe D., Eraus S., Woodworth J. (1986): Heavy metal toxicity to fish and the influence of water hardness. Arch. Environ. Contam. Toxicol, 15,
- Patel S.K. and P.G. Parmer, (1993) : Studies on acute *Heteropneustes fossilis* (Bloch). Dhaka University Stud. toxicity of endosulfan on protein contents of Bangladesh, 26: 115-112.
- Patnail B.B. , K. A. Sudhan and M. Selvanayagam. (2002) : Toxicity of two finger lings of a carp *Sarotherodon mossambicus* by a synthetic pyrethroid fenvalerate .Ind. J. Environ. Ecoplan. 6: 551-558.

- Peakall D., (1992) : Animal Biomarkers as Pollution Indicators. Chapman & Hall, London. 1992.
- Poleksic V., & Mitrovic-Tutundzic, V. In Müller, R., & Lloyd, R (Eds.). (1994). Fish gills as a monitor of sublethal and chronic effects of pollution. *Sublethal and Chronic Effects of Pollutants on Freshwater Fish* (pp.339-352).
- Prakash Chandra koal., Mastan S. A., and t. A. Qureshi (2007) : Sublethal effect of Dichlorvos (DDVP) on certain biochemical parameters of *Channa gachua* (Ham). Journal of Herbal Medicine and Toxicology. 1 (2) 29-32.
- Pulla Rao, U.D. V.P., and K.D., Mishra 1997: acute toxicity of herbicides atrazine to fish, *Puntius ticto*: A comparison of bioassay test methods. J. Environ. Polln. 4; 161-165.
- Pundir Rekha, Saxena A.B., (1991) : Studies on pesticide toxicity to fresh water fish *Channa punctatus*. *J Hydrobio*, 7 (1) (1991), 61-63 [10 Ref].
- Radha G.S., Logaswami and Logankumar, (2005): Sub lethal toxicity of dimethoate on protein, glucose and cholesterol contains in the fish *Cyprinus carpio*. Nat. Environ. Polln. Tech. 4(2) : 307-310.
- Radhakrishniah K, Venkataramana P, Suresh A, Sivaramkrishna B (1992): Effects of lethal and sublethal concentrations of copper on glycolysis in liver and muscle of the freshwater teleost, *Labeo rohita* (Hamilton). *J Environ Bio*, 13 (1) (1992), 63-68 [23 Ref].
- Rahman M.Z., Hossain Z., Mullah MFR, Ahmed GU (2002): Effect of Diazinon 60EC on *Anabus testudineus*, *Channa punctatus* and *Barbades gomnotus*. NAGA. The ICLARM Quarterly, 25: 8-11.
- Rainza-Paiva, M.J.T., Ishikawa, C.M., Das Eiras, A.A.; Felizardo, N.N., (2000) : Haematological analysis of Chara, *Pseudoplatystoma fasciatum* in captivity. European Aquaculture Soc. Special Pub. Vol. 28, 590 pp.
- Raizada M.N., K.K., Jain and Raizada S., (1983) : Monthly variation in the haematocrit values (PCV) in a teleost, *Cirrhinus mrigala* (Ham). Comp. Physiol. Rcol., 8: 190-196.
- Rajamannar K., and L., Manohar (1998) : Effect of pesticides on oxygen consumption of fresh water fish *Lebio rohita*. J. Ecobiol. 10: 205-208.
- Rajendranath T., and Benerji. G., (1991) : Toxic effect of dichlorovos (DDVP) on the gills of fresh water teleost *Clarias batrachus* (Linn). Environ. Ser. V. 185-191.

- Rajeshwari K.S.J., Reddy D.C., Reddy and R., Ramamurthy (1989) : Effect of dichlorvos on certain blood parameters of the fish, *Clarias batrachus*. Environ. Ecol. 7 : 933-934.
- Ramawamy M., and G.T., Reddy, (1988) : A Comparative study of haematology of three air-breathing fishes. Proc. Indian Acad. Sci., 12: 381-385.
- Ramesh M., and Saravanan M., (2008) : Haematological and biological responses in fresh water fish *Cyprinus carpio* exposed to Chlorpyrifos. International Journal of Intrigative Biology. IJIB, 853-863.
- Ramesh M., Manavalaramanujam Sivakumari K. (1992) : Effect of water hardness and the toxicity of malathion on haematological parameters of the freshwater fish *Cyprinus carpio*. *J Ecotoxicol Environ Monit*, 2 (1) (1992), 31-34 [10 Ref].
- Rand G.M. (1977) : The effect of exposure to a subacute concentration of parathion on the general locomotor behavior of the goldfish. Bull. Environ. Contam. Toxicol., 18(2), 259-266, (1977)
- Rand G.M., and Petrocelli S.R., (1985) : Introduction. In: Fundamentals of Aquatic Toxicology : Methods and Applications. (Edited by : G.M. Rand and S.R. Petrocelli), Hemisphere Publishing Corporation, USA, pp. 1 – 8.
- Rand S.M., and S.R., Petrocelli., Johnson, C.M., and M.C.F., Toledo (1993) : Acute toxicity of endosulfan to the fish *Hypessbrycon bifasciatus*. Hemisphere Publishing Cooperation, Washington, pp: 666.
- Ranke B., and Rybicki (1975) : Effect of organophosphorus pesticides on aquatic organism. Pocz. Panst. Zakl, Hig. 26. 293-300.
- Ransford K. D., (1978) : Toxicity of organophosphorus insecticides: Comparison to the toxicity of metabolites with parent compounds using an intracerebral injection method. Pesticides, Biochemistry and Physiology. (8) 302-316.
- Rashed M. N., (2001) : Cadmium and lead levels in fish (*Tilapia nilotica*) tissues as biological indicator for lake water pollution. Environ. Monit. Assess, 68: 75-89.
- Rath S., and Mishra B.N., (1980) : Age related changes in oxygen consumption by the gill, brain and muscle tissue of *Tilapia mossambica* (Peters) exposed to Dichlorvos (DDVP). Environ. Poll. (23) 96-104
- Rathore H.S., and Kothari S.K., (1986) : Adaptive respiratory behaviour of catfish exposed to CdCl₂ under experimental conditions. Poll. Res., 5(3&4), 129-146.

- Ravindra Nath and Benerjee V., (1999) : Influence of lethal and sublethal of roger on the blood parameters of fresh water fish *Heteropneustes fossilis*. Environmental and Ecology 17 (4) 940-944.
- Reddy D.C., P., Vijaykumari V., Kalarani and R. W., Davis (1992) : Change in erythropoietic activity of *Sarothodon mossambicus*. Exposed to sub lethal concentration of the herbicides, Diuron. Bull. Environ. Contam. Toxicol: 49: 730-737.
- Rehulka J. (2002a) : Aeromonas causes severe skin lesions in rainbow trout (*Oncorhynchus mykiss*): Clinical pathology, haematology and biochemistry. Acta Vet Brno, 71: 351-360.
- Rehulka J. (2002b): Effect of polychlorinated biphenyls Delor 103 on some haematological and biochemical indices of the blood plasma of the rainbow trout, *Oncorhynchus mykiss* (Walbaum). In: The second PCB workshop. Recent Advances in the Environmental Toxicology and Health Effects of PCBs. May 07-11, 2002. Brno, Czech Republic. (Book of Abstracts): 36.
- Reid S. D., Mc Donald, D. G., (1988) : Effects of cadmium, copper and low pH on ion fluxes in the rainbow trout, *Salmo gairdneri*. Can. J. Fish. Aquat. Sci, 45: 244 – 253.
- Reitmayer J, Felzl L, Roth Z, Tich.M. (2006) : Distribution of components of binary mixtures between aqueous and gaseous phases. *Chem Listy* 100: 42-47.
- Risbourg S.B., and J. Bastid, (1995) : Hepatic perturbations induced by a herbicide (atrazine) in juvenile grey mullet *Liza ramada* (Mugilidae Telostei): an ultra structural study. *Aquat Toxicol.*, 31: 217-29.
- Roberts, R.O. and S.G. Berk. (1993) : Effect of copper, herbicides, and a mixed effluent on chemoattraction of *Tetrahymena pyriformis*. *Environmental Toxicology and Water Quality* 8:73-85.
- Rogers., J.T., Richards, J.G. and Wood, C.M. (2003) : Ionoregulatory disruption as the acute toxic mechanism for lead in the rainbow trout. *Aquatic Toxicol.*, 64 (2): 215-34.
- Rojik J., Nemcsok J, Borross L (1983): Morphological and biochemical studies on liver, kidney and gill of fish affected by pesticides. *Acta Biol Hungarica* 34: 81-92
- Roopa Methew., M.K., Kanagaraj and R. Manavalaramanjuam (1997) : Lead nitrate toxicity on ventilation frequency, oxygen consumption and haemoglobin content in fish *Cyprinus carpio*. *Poll Res.* 16 (1) : 51-53.
- Roux D.J., (1994): Role of various contaminants in water quality assessment and a case study on the Kokolite river, *Eastern Transvaal*.

Doctral thesis, RAU, 130 pp.

- Roy, D.N., S. K. Konar, S. Banerjee, D. A. Charles, D. G. Thompson, and R. Prasad. 1989b. Uptake and persistence of the herbicide glyphosate in fruit of wild blueberry and red raspberry. *Journal of Forest Research* 19:842-847.
- Roy, D.N., S. K. Konar, S. Banerjee, D. A. Charles, D. G. Thompson, and R. Rueppel ML, Brightwell BB, Schaefer J, and Marvel JT, (1977) : Metabolism and degradation of glyphosate in soil and water. *J. Agric. Food Chem.* 25:517-528.
- Rueppel, M.L., B.B. Brightwell, J. Schaefer and J.T. Marvel. (1977): Metabolism and degradation of glyphosate in soil and water. *Journal of Agricultural and Food Chemistry* 25:517-528.
- Ruparelia S.G., Verma Yogendra, Saiyed SR, Rawal UM (1991) : Changes in differential white blood cell (WBC) count in fish, *Sarotherodon mossambica* (Peters) under prolonged exposure to cadmium. *Indian J Environ Toxicol*, 1(1) (1991), 59-64 [21 Ref].
- Sadhu D.N., (1993) : Toxicity of organophosphorous insecticides monocil to air breathing fish *Channa punctatus* (Bloch) .*J. Ecotox. Environ. Mon.* 3: 133-136.
- Sadhu H. S., Gupta A. R., and Komble G.S., (2001) : The toxicological study of Dusan 50 fish *Channa gachua* 9 (2) : 11-13.
- Sailajaraghuram D. and B.N., Naidu (1989) : Hematology in fresh water fish *Tilapia mossambica* (Pester) with special reference to methyl parathion toxicity. *Nat. Acad. Sci. Lett.* 12: 77-79.
- Salim Mustfa and Ajmal Murad (1984) : Servival, beheviour response and haematological profile of cat fish *Heteropneustes fossilis* exposed to DDT. Vol. 31, No. 1 (1984).
- Samprath K, Velamnial S, Kennedy IJ, James R (1993) : Haematological changes and their recovery in *Oreochromis mossambicus* as a function of exposure period and sublethal levels of Ekalus. *Acta Hydrobiol.* 35: 73-83.
- Santha kumar M., M. Balaji and K. Ramuudu (2000) : Histopathological lesion in the brain of teleost *Anabas testudineus* exposed to monocrotophos. *J. Enverion. Biol.* 21: 197-199.
- Santillo D.J., D. M. Leslie Jr., and P. W. Brown. (1989a): Response of small mammals and habitat to glyphosate application on clearcuts. *Journal of Wildlife Management* 53(1):164-172.
- Santillo D.J., P. W. Brown, and D. M. Leslie, Jr.. (1989b): Response of songbirds to glyphosateinduced habitat changes on clearcuts. *Journal*

of Wildlife Management 53(1):64-71.

- Sarah J., Stoughton Karsten, Liber Joseph, Allan Cessna (2007) : Acute and chronic toxicity of Imidacloprid to the aquatic invertebrate *Chironomus tentans* and *Hyaella azteca* under constant and pulse exposure conditions. Arch Environ Contam Toxicol . 54: 662-673.
- Sarkar (1990a) : Acute toxicity of 2, 4- D to fish effects of temperature. Proc. Zool. Soc. 43: 45-48.
- Sarkar (1990b) : Acute toxicity of 2, 4- D to *Cyprinus carpio*. Environ. Ecol. 8 : 316-318.
- Sarkar (1997b) : Toxicity of simazine, DAP and rock phosphate to fish at various water temperature. Geob. 24: 142-145.
- Sastry A. and R. A. Agrawal. (1993) : Effect of cypermethrin on lactate dehydrogenase and cytochrome oxidase of fish. Bull. Environ. Contam. Toxicol. 51: 445- 452.
- Sastry and Sunita. (1982) : Effect of cadmium chromium on the intestinal absorption of glucose in snake head fish *Channa punctatus*. Toxicology Letters, 10: 293-296.
- Sastry K.V and K. Sharma (1981) : Diazinon induced histopathological and haematological alterations in a freshwater teleost, *Ophiocephalus punctatus*. Ecotoxicology Environment Safety, 5: 329-340.
- Sastry K.V. and A. Gupta (1994) : Hematological study on the effect of cadmium and di-methoate alone and in combination on a fresh water teleost fish *Channa punctatus* . J. Environ. Polln. 1: 133-139.
- Sastry K.V., Siddiqui A.A. and Samuel. M. (1988) : Acute and chronic toxic effects of the carbamate pesticide sevin on some hematological Biochemical and enzymatic parameters in the fresh water teleost fish, *Channa Punctatus*. Acta Hydrochim. Hydrobiol. 16 (6) : 625-631.
- Sastry N.N., V.K. Das and A. K., Shrivastava. (1997) : Formation and propoxur induced ionic imbalance and sub lethal deformity in a cat fish *Heteropneustes fossilis*. J. Environ. Biol. 18: 157-163.
- Saufy H., M. K., Soliman E.M., El-Manakhly and A. Y., Gaafar (2007) : The biochemical and pathological investigations on *Monex tilapia* following chronic exposure to caebofuran. Global Veterinaria 1 (1) : 45-52.
- Sawada Y., Nagai Y., (1987) : Roundup poisoning and its clinical observation: Possible involvement of surfactant. *J Clin Exp Med* 143:25-27, 1987.
- Schaefer A. A., (1925): The number of red corpuscles in in fish's relation

- to starvation and seasonal cycle. J.gen. Physiol., Vol. 7, pp. 341-343.
- Scheck C. B., (1990): Physiological, behavioural, and performance indicators of stress. In: Adams S.M. (eds) Biological indicators of stress in fish, American Fisheries Society, Bethesda, MD pp 29 - 37.
- Schulz-Jander, D. A., Leimkuehler, W. M. and Casida, J. E. (2002): Neonicotinoid Insecticides: reduction and cleavage of imidacloprid nitroimine substituent by liver microsomal and cytosolic enzymes. Chem. Res. Toxicol. 9:1158-65.
- Selvakumar S., Ajmal Khan S, Kumaraguru A. K., (1996) : Acute toxicity of some heavy metals, pesticides and water soluble fractions of diesel oil to the larvae of some brachyuran crabs. *J Environ Bio*, 17(3) (1996), 221-226 [25 Ref].
- Servizi J. A., R. W. Gordon, and D. W. Martens. (1987): Acute toxicity of Garlon 4 and Roundup herbicides to salmon, *Daphnia*, and trout. Bull. Environ. Contam. Toxicol. 39:15-22.
- Servizi J.A., Gordon R.W., Martens D.W. (1987) : Acute toxicity of Garlon 4 and Roundup herbicides to salmon, *Daphnia* and trout. Bull Environ Contam Toxicol 39: 15-22.
- Sexena V. S., Trivedi and D. N., Saksena (1997) : Acute toxicity of two organophosphorous pesticides nuvan and dimecron to a fresh water murrel, *Channa orientalis* (Schneider) J. Environ. Polln. 4 : 79- 83.
- Shah S.L., (2006) : Haematological parameters in tench, *Tinca tinca* after short term exposure to lead. Journal of applied toxicology, 26 (3): 223-228.
- Shaikha Y.A., and P.K., Hiradher (1985) : Fluoride induced changes in blood Glucose, tissue glycogen and succinate dehydrogenase (SDH) Activity in the mudskipper *Beleophthalmus dessoignei*. Proc. Symp. Assess. Envir. Pollu, 93-99.
- Sharma R. C., Gupta N (1984): Carbon tetrachloride induced haematological alterations in *Clarias batrachus* (L.) J. Environ. Biol. 3: 127-131.
- Shastri K.V., and Siddiqui A. A. (1982) : Chronic toxic effect of carbamate pesticides sevin on carbohydrate metabolism in snake head fish *Channa punctatus*. Toxicology letters, 14, 123-130.
- Shastri K.V., Abad. Siddiqui. A. (1984): Some hematological, biochemical and parameters of fresh water teleost fish (*Channa punctatus*) exposed sub lethal concentration of quinaldine. pest.Biochem.Physiol,22:8-13.
- Shastri K.V., Sachdeva S., Rathee P. (1997): Bioaccumulation of cadmium

- and copper and their combination of enzymological and biochemical parameters in *Channa punctatus*. *J. Env. Biol.* 18(3), 291-303.
- Sheets L.P. (1994b): An Acute Oral Neurotoxicity Screening Study with Technical Grade Imidacloprid (NTN 33893) in various fishes . Miles Inc. (Mobay). Study No. 106348. DPR Vol. 519500473.
- Sheets L.P. (2001) An Developmental Neurotoxicity Study with Technical Grade Imidacloprid in different fishes. Bayer Corporation, Agricultural Division Toxicology; Kansas, USA. Study No.110245. DPR Vol. 51950-0474 # 209393.
- Shekhar P., and I., Christy (1996) : Hematological changes in the fresh water cat fish (*Mystus vitattus*) exposed to sublethal concentration of Phosphamidon. *J. Ecol.* 8(1) : 25-28.
- Sherry P. M., and Abidi Z. Z., (2002) : Erythrocyte and Leucocyte Response to Lead and Cadmium poisoning in fresh water teleost *Channa punctatus* (Bloch). Proceedings 89th session of Indian Science Congress Zoology section ABS No. 171. University of Lucknow.
- Shobha R.J.V., Venketeswarlu P.V., and Jainih C. (1989) : Change in carbohydrate metabolism in fresh water fish *Heteropneustes fossilis* . *Ecotoxicolgy and environmental Safety* (5) 412-417.
- Shrivastva A. K. and K. A., Shrivastava (1987) : Acute toxicity of kepon to the fresh water fish *Heteropneustes fossilis* . *Nat. Acad Sci. Lettters.* 10: 439-441.
- Shrivastva A. K. and N. N., Shrivastva (1987) : Hematological changes in fresh water fish *Heteropneustes fossilis* following exposure to aldrin. *J. Nat. Con.* (1) : 85-88.
- Shrivastva A. K., and Singh N. N., (1981) : Effect of aldrin on carbohydrate metabolism in indian cat fish (*Heteropneustes fossilis*) *Pharmacol. Ex. Toxicol.* 49. 266-269.
- Shukla V., and Shastry K.V. (1998) : Acute Toxic Effects of Cadmium on various parameters in the fresh water fish. *Proc. Acad. Environ. Biol.*, 7(2), 155-160.
- Siddiqui A.Q., and S. M., Nasim (1989) : The hematology of mrigal, *Cirrhina mrigala* (Ham.) Teleostei: Cyprinidae. *Anat. Anz.*, 146: 262-269.
- Sievers G., P. Palacios, R. Inostroza and H. Doelz (1995) : Evaluation of the toxicity of 8 insecticides in *Salmo Salar* and the *in-vitro* effects against the Isopade parasite, *Ceratomyxa gandichandi* *Aquacul.*, 134: 9-16.

- Sikoki F. D. & Enajekpo H.O.S. (1991) : Effects of the water soluble fraction of crude oil on some haematological factors and liver glucose in *O. niloticus*. Paper presented at the 6th annual conference of the Nigerian Association for aquatic science held at the University of Benin, 11 pp.
- Silbergeld E.K., (1974) : Blood glucose : A sensitive indicator of environmental stress in fish. *Bull. Environ. Contam. Toxicol.* 11: 20-25.
- Singh B. B., & Narain A.D. (1982) : Acute toxicity of thiodan to catfish, *Heteropneustes fossilis*. *Bull. Environ. Contam. Toxicol.*, 28:22-27.
- Singh B., and Bhati DPS., (1991) : Effect of Zinc chloride on the morphology of blood in *Channa punctatus* (Bloch) *Natural Environment*, 8; 27-32.
- Singh H. H., Srivastava A. K., (1982) : Effect of formothion on carbohydrate metabolism in Indian catfish (*Heteropneustes fossilis*). *Environ. Res.* 28: 335-339.
- Singh H.S., Reddy T.V., (1990) : Effect of copper sulphate on hematology, blood chemistry and hepatosomatic index of an indian catfish, *Heteropneustes fossilis* (Bloch) and its recovery. *Ecotoxicol Environ Saf* 20: 30-35
- Singh N. N., Srivastava, K. A. & Srivastava K. A., (1992) : Blood dyscrasis in fresh water Indian cat fish after acute exposure to a sublethal concentration of propoxur. *ACTA Hydrobiol.*, 34: 189-195.
- Singh S and D. N., Sadhu (2001) : Toxicity and behavior of rogor (dimethoate) exposed to *Channa punctatus* (bloch) .J. *Environ. Polln.* 8: 377-378.
- Singh Snehlata, Sadhu D. N., (2001) : Toxicity and behaviour of Rogor (dimethoate) exposed *Channa punctatus* (Bloch). *J Env Polln*, 8(4) (2001), 377-378 [8 Ref].
- Sinha Manoj Kumar, Datta Munshi Jayashree (1994) : Toxicity of alkaloid extract of *Datura meli* L. on *Heteropneustes fossilis* (Bloch) and *Anabas testudineus* (Bloch) in relation to behaviour of fishes. *J Fresh water Bio*, 6(1) 79-85 [20 Ref].
- Soivio A., Westman, K. & Nyohal, K., (1974) : Changes in the haematocrit values in blood samples treated with and without oxygen: a comparative study with four salmonid species. *J. Fish Biol.* 5, 763-769.
- Song M.Y., Stark J.D. and Brown J.J. (1997) : Comparative toxicity of four insecticides, including imidacloprid and tebufenozide, to four aquatic arthropods. *Environ Toxicol Chem* 16:2494-2500.

- Sreenivasulu C., R., Samsan and K., Jayantha Rao. (1991) : Change in chromatophores and amino acid pool in *Tilapia mossambica* exposed to dimecron. *Environ. Ecol.* 13: 626-628.
- Srivastava A. K., (1981) : Effects of acute exposure of methyl parathion on carbohydrate metabolism of Indian catfish (*Heteropneustes fossilis*). *Acta Pharmacol. Toxicol.* 48: 26-34
- Srivastava R.K. and Srivastava S., (2002) : Cadmium (Cd) induced hematological changes in a freshwater catfish, *Heteropneustes fossilis*. *J. Adv. Zool.*, 23(1), 23-29.
- Stein J. E., Collier, W. L., Reichert, E, Casillas, T. and Hom T (1992) : Bioindicators of contaminant exposure and sublethal effects: Studies with benthic fish in Puget Sound, Washington. *Environ. Toxicol. Chem.* 11: 701 – 714.
- Stine K.E., and T.M. Brown, (1996) : Principles of Toxicology. CRC press, Inc. USA., pp: 211-225.
- Stout B.A., (1976) : Energy for World agriculture, FAO. Agric. Sem. 7:286 pp
- Subramanian S. and Ramalingam K. (2003) : Effect of DDT, malathion and mercury on the haematology of *Sarotherodon mossambicus*. *J. Ecotoxicol. Environ. Monit.*, 13(2), 159-160.
- Suege J., and Bluzat, R., (1983) : Chronic toxicity of three insecticides, Carbaryl, Fenthion and Lindane, 106(1):65-72
- Sullivan T. P. and D. S. Sullivan. (1979): The effects of glyphosate herbicide on food preference and consumption in black-tailed deer. *Can. J. Zool.* 57:1406-1412.
- Svoboda, M; Luskova, V; Drastichova. Zlabek, V, (2001) : The effect of diazinon on haematological indices of Common Carp (*Cyprinus carpio L.*). *Acta vet. BRNO*, 70: 457-465
- Swif, D.J. and R. Lloyd. (1974) : Changes in urine flow rate and hematocrit value of rainbow trout *Salmo gairdneri* (Richardson) exposed to hypoxia. *Journal of Fisheries Biology* 6:379-387
- Tai T., Yamashita M, Wakimori H., (1990) : Hemodynamic effects of Roundup, glyphosate and surfactant in fishes. *Jpn J Toxicol* 3:35-88.
- Talbot A.R., Shiaw M-H, Huang J-S, (1991) : Acute poisoning with a glyphosate-surfactant herbicide (Roundup): A review of 93 cases. *Hum Exp Toxicol* 10:1-8, 1991.
- Tavares-Dias, M., F. R. Morsaes (2004) : Haematology in teleosts fish. M. Tavares-Dias (ed.),
- Terry G., (1987) : Breaking the pesticides Habit. 18. Viveen, W.J.A.R.,

- C.J.J. Rhcher, P.G.W.J. Van Oordt, Alternatives to 12 hazardous Pesticides. IASA publication.
- Thakur G. K., and P. K. Pandey (1991) : BHC (Gammexene) Poisoning effect on leucocytes of an air breathing fish, *Clarius batrachus* (Linn). J. Environ. Biol. 11(2), 105-110.
- Thielen K.D., E.P. Jackson and D. Penner. (1995a): The basis for the hard-water antagonism of glyphosate activity. *Weed Science* 43:541-548.
- Tilak K.S., Veeraiah K, Anita Susan T, Yacobu K (2001) : Toxicity and residue studies of fenvalerate to some selected freshwater fishes. *J Environ Bio*, 22(3), 177-180.
- Tilik K.S., K. veeraiah and D. K. Rao (2005) : The effect of chlorpyrifos, an organophosphate in acetyl choline esterase activity in fresh water fishes. *J. Environ. Biol.* 26 (1) : 73-78.
- Tiwari C. (1995) : Laboratory monitoring of fungicides toxicity to fresh water fish *Puntius ticto* , Ph. D. thesies, Barkatullah university, Bhopal.
- Tominack R.L., Yang G-Y, Tsai W-J: (1991) : Taiwan National Poison Center survey of glyphosate-surfactant herbicide ingestions. *Clin Toxicol* 29:91-109, 1991.
- Tomizawa M. and Casida, J. E. (2000): Imidacloprid, thiacloprid, and their imine derivatives upregulate the alpha 4 beta 2 nicotinic acetylcholine receptor in M10 cells. *Toxicol. Appl. Pharmacol.* 169:114-20.
- Tomlin C.D.S. (1997) : The pesticide manual: Incorporating the agrochemicals handbook, 10th ed. British Crop Protection Council, Farnham, Surrey, UK.
- Tomlin C.D.S. (2000) : The Pesticide Manual, 12th Edition. British Crop Protection Council, Surrey, United Kingdom.
- Tort L., Torres P (1988). The sublethal concentration of cadmium on haematological parameters in dogfish. *Squalorhina canicula*. *J. Fish Biol.* 32: 277-282.
- Tripathi G. and S. P., Shukla. (1988) : Toxicity bioassay of technical and commercial formulation of carbaryl to the fresh water catfish *Clarias batrachus*, *Eco tox. Saf.* 15(3) 277-281.
- Trivedi S., and D. N. Saksena. (1999) : Acute toxicity and behavioral responses of an organophosphorus pesticides , nuvan (DDVP) to fresh water fish *Clarias batrachus*, (Linnaechus) *J. Environ. polln.* 6: 53-57.
- Tsuda T., Kojima M., Harda A., Nakajima and Akoki S., (1997) : Acute

toxicity of, accumulation and excretion of organophosphorus insecticides and their oxidation products in kill fish Chemosphere. (35) 939-949.

United States Environmental Protection Agency (USEPA). (1993): Memo regarding the Risk to Human Health From Exposure to Pesticide. Imidacloprid. March 4, 2003. DPR Vol. 51950-0456

United States Environmental Protection Agency (USEPA). (1994): Rules and Regulations. Imidacloprid; tolerances for residues. Federal Register Vol. 63, No. 57 / March 21 1994.

Univ and Burdwan (1999) : Effect of cadmium on hematological parameters in young and adult fish *Labeo rohita* (Hamilton). *Env Eco*, 17(4), 999-1004.

USDA (1984) : Herbicide Background Statement: Glyphosate. In: Pesticide Background Statements. Vol 1: Herbicides USDA Forest Service, Agricultural Handbook No 633: G-1-72.

USEPA (1993b) : A Review of Ecological Assessment Case Studies from a Risk Assessment Perspective. EPA/630/R-92/005.

USEPA, (1993a): Re-registration Eligibility Decision (RED), Glyphosate. Office of Prevention, Pesticides and Toxic Substances, U.S. Environmental Protection Agency, Washington, DC. 738-F-93-011. (738-R-93-014).

Vaala and Mitchell (1970) : Blood oxygen tension changed in acid exposed brook trout. *Proc. Pac. Acad. Sci.* 44, 41-44.

Valicre E.J., and C.J. Stickney (1989) : Hypoxia in fishes. Uni. of Cicago. London, 1989.

Van Vuren, J.H.J. (1986) : The effects of toxicants on the haematology of *Labeo umbratus* (Teleostei; cyprinidae). *Comparative Biochemistry and Physiology* 83C: 155-159.

Vander Oost, R., Beyer J., and Vermeulen, N.P.E. (2003) : Fish bioaccumulation and biomarkers in environmental risk assessment: a review. *Environ. Toxicol. Pharmacol.*, 13: 57-69.

Vatukuru S.S., (2005) : Acute effects of hexavalent chromium on survival, oxygen consumption, haematological parameters and some biological profiles of indian major carp, *Lebio rohita*.

Vdesek J., Z. Svoboda, V., Piackova (2009) : Effect of acute exposure to bifenthrin on some haematological, biochemical and histopathological parameters of rainbow trout (*Oncorhynchus mykiss*). *Veterinari Medicina* 54, 3. 13-137.

Verma G.P. and Pranamita Panighari (1998): Effect of agrofen on blood

- parameters of *Oreochromis mossambica*. Proc. Nat. Acad. Sci. India. 68. (B) 1,11-18.
- Verma S.R., Bansal A.K., Gupta N., Pal A.K., Tyagi A.K., Bhatnagar V. Kumar and R.C. Dalela. (1979): Acute toxicity of twenty three pesticides to fresh water teleost, *Sacbranchus fossilis*. Proc. Symp. Environ. Biol. 481-497.
- Verma S.R., Bansal S.K, Gupta AK, Dalela R.C., (1979): Pesticide induced haematological alterations in a freshwater fish. *Sacchbranchus fossiles*. Bull. Environ. Contam. Toxicol. 22: 467-474.
- Verma S.R., S., Rani S.K., Bansal and R.C., Dalela (1981) : Evolution of three comparative toxicity of thiox, dichlorovas and carbofuron to fresh water teleost , *Ophiocephelus punctatus* and *Mittus vittatus*. Acta. Hydrochim. Hydrobiol. 9: 119-129.
- Verma S.R., S.K., Bansal A.K.,Gupta N., Pal A.K., Tyagi A.K., Bhatnagar V., Kumar and R.C., Dalela (1990) : Bioassay with twenty three pesticides to fresh water teleost, *Sacbranchus fossilis*. Wat. Res. 16: 525-530.
- Vivekanandan E., Pandian T.J., (1977) : Surfacing activity and food utilization in a tropical air-breathing fish exposed to different temperatures. *Hydrobiologia* 1977; 54: 145-160.
- Vosyliene M.Z., (1999) : The effect of heavy metals on Haematological indices of fish Acta Zoologica Lituanica 9 (2): 76-82.
- Voyer R.A., (1985) : Effect of dissolved oxygen concentration on acute toxicity of cadmium to the Mummicho Fundulus heteaoclitus (L.) at various salinities. *Trans. Am. Fish Soc.*, 104, 129-134.
- Vutukuru S.S., (2000) : Acute Effects of hexavalent chromium on survival, oxygen consumption, haematological parameters and some biochemical profiles of the Indian major Carp, *Labeo rohita* Int. J. Environ. Res. Public Health, 2005, 2(3), 456-462.
- Wademeyer and Yasutake (1977) : A clinical methods for the assessment of the effect of environmental stress on fish health. Technical papers of the U.S. Fish and Wildlife Service, No. 89,14pp.
- Wannee Jiraungkoorskula E., Suchart Upathama B., Maleeya Kruatrachuea, Somphong Sahaphongc, Suksiri Vichasri-Gramsas and Prayad Pokethitiyooka (2002) : Histopathological effects of Roundup, a Glyphosate herbicide, on Nile tilapia (*Oreochromis niloticus*) *ScienceAsia* 28 : 121-127.
- Warren M. L., and B. M., Burr. (1994) : Status of freshwater fishes of the United States: overview of an imperiled fauna. *Fisheries* 19(1):6-18.

- Washington D.C., Weber, D.N., Spieler R. E., (1994) : Behavioural mechanisms of metal toxicity in fishes. In: Malins D.C; Osstrander, G.K. (eds) Aquatic Toxicology, molecular, biochemical and cellular perspectives, Lewis , Boca,raton, FL, USA, pp421 – 467.
- Watson W.J., K.L., Brown and M.J., Lydy (1997) : Diurnal fluctuations in toxicity to fish species *Gambusia affinis* and *Notropis lundibundis*. Bull. Environ. Contam. Toxicol. 59: 414-421.
- Weber D.N., Spieler R. E., (1994) : Behavioural mechanisms of metal toxicity in fishes, Aquatic Toxicology, molecular, biochemical and cellular perspectives, Lewis , Boca,raton, FL, USA, pp421 – 467.
- Wedemayer G. A., Yasutake, W. T. , (1987) : Clinical methods for the assessment of the effects of environmental stress on fish health. *US Fish and wildlife service Tech. Paper* 89.
- Wedemeyer G., D. J. Mcleay, and C. P., Good year (1984) : Assessing the tolerance of fish and fish population to environmental stress: The contaminant effects on fisheries Eds. W. V Cairns, P. V. Hodson and J.O Nriagu 184-195.
- Wedemeyer G.A., Macley D.J., and Goodyear C.P., (1976) : Assessing the tolerance of fish and fish population to environmental stress : The problems and methods of monitoring, p. 161-165. . John Wiley and sons, New York.
- Weis J. S., Smith, G., Zhou, T., Santiago-Bass, C., Weis, P. (2001) : Effects of Contaminants on Behavior: Biochemical Mechanisms and Ecological Consequences. *Bio Science* 51(3):209-217.
- Wepener V., (1997) : Metal Ecotoxicology of the Olifants River in the Kruger National Park and the Effect thereof on Fish Haematology. PhD-Thesis, RAU.
- Wesseling C., Castillo L, Elinder C.C., (1993) : Pesticide poisonings in Costa Rica. *Scand J Work Environ Health* 19:227–235, 1993.
- Westlake G.F., Kleerekoper G.H., and Matis J., (1974): The locomotor response of goldfish to a steep gradient of copper ions. *Water Resources Research*. 10:103-105.
- Williams G.M, Kroes R., and Munro I.C. (2000): Safety Evaluation and Risk Assessment of the Herbicide Roundup and Its Active Ingredient, Glyphosate, for Humans. *Regulatory Toxicology and Pharmacology* 31, 117-165.
- Wilson R.W. and Taylor, E.W. (1993): The physiological responses of freshwater rainbow trout, *Onchorynchus mykiss*, during acute exposure. *J. Comp. Physiol.* 163b: 38-47.

- Wilson R.W., Taylor E.W. (1993) : The physiological responses of freshwater rainbow trout, *Onchorynchus mykiss*, during acute exposure. J. Comp. Physiol., 163b: 38-47.
- Yamamoto K-I, Itazawa Y., and H., Kobayashi. (1980): Supply of erythrocytes into the circulating blood from the spleen of exercised fish. Comparative Biochemistry and Physiology 65A:5-13.
- Yamamoto K-I., Itazawa Y., and H. Kobayashi. (1983): Erythrocyte supply from the spleen and hemoconcentration in hypoxic yellowtail. Marine Biology 73:221-226.

ABBREVIATIONS AND SYMBOLS

1. < smaller than
2. = equal to
3. μg Micro gram
4. Ach acetylcholine
5. AChE acetylcholine enzyme
6. BOD Biological oxygen demand
7. Cas catecholamine
8. cm centimeter
9. CNS central nervous system
10. DO dissolved oxygen
11. ESR Erythrocytes sedimentation rate
12. FAO Food and Agriculture Organization of the United Nations
13. Hb% Haemoglobin percentage
14. L liter
15. LC_{50} lethal concentration, 50% kill
16. MCH Mean Corpuscular Haemoglobin
17. MCHC Mean corpuscle haemoglobin concentration
18. MCV Mean Corpuscular Volume
19. ml milliliter
20. MW molecular weight
21. nAChR Nicotinic acetylcholine receptor
22. ODM Oxydemeton-methyl
23. PCV Packed cells volume
24. PEP phosphoenolpyruvate
25. ppm parts per million

- 26. r.p.m. Revolution per minute
- 27. RBC red blood cells
- 28. TEC Total erythrocytes count
- 29. TLC Total leucocytes count
- 30. Unit (blood) (mg/dl)
- 31. WHO World Health Organization
- 32. viz videlicet (namely)
- 33. i.e. id est (that is)
- 34. a.e. acid equivalents
- 35. a.i. active ingredient

ANNEXURE

Publications

- K. Zahra and Saurabh Shreshth (2005): Variation in blood parameters due to Glyphosate on fresh water fishes *Heteropneustes fossilis*. 16Th All India congress of Zoology and national symposium recent advance in animal research with special emphasis on invertebrates. October pp: 96.
- K. Zahra and Saurabh Shreshth (2005): Haematological effects in *Channa punctatus* after acute and sub acute exposure of Glyphosate. 16Th All India congress of Zoology and national seminar. December 07: PS 361.
- K. Zahra and Saurabh Shreshth (2006): Effect of glyphosate on various blood parameters of fresh water fishes, *Heteropneustes fossilis*. Flora and fauna Vol. 12 No. 1 pp 100-104.